

TEXTE

XX/2022

Review of internationally proposed critical levels for ammonia

Proceedings of an Expert Workshop held in Dessau and
online on 28/29 March 2022

TEXTE XX/2022

Ressortforschungsplan of the Federal Ministry for the
Environment, Nature Conservation and Nuclear Safety

Project No. (FKZ) 3718 63 2010

Report No. (UBA-FB) XXX

Review of internationally proposed critical levels for ammonia

Proceedings of an Expert Workshop held in Dessau and
online on 28/29 March 2022

prepared by

Jürgen Franzaring and Julia Köster
Institute of Landscape and Plant Ecology,
University of Hohenheim, Stuttgart

On behalf of the German Environment Agency

Imprint

Publisher

Umweltbundesamt
Wörlitzer Platz 1
06844 Dessau-Roßlau
Tel: +49 340-2103-0
Fax: +49 340-2103-2285
buergerservice@uba.de
Internet: www.umweltbundesamt.de

[f/umweltbundesamt.de](https://www.facebook.com/umweltbundesamt.de)

[t/umweltbundesamt](https://twitter.com/umweltbundesamt)

Report prepared by:

University of Hohenheim Institute of Landscape and Plant Ecology
Otilie-Zeller Weg 2
70599 Stuttgart
Germany

Report completed in:

August 2022

Edited by:

Section II 4.3 Air Quality and Terrestrial Ecosystems
Markus Geupel (Scientific Support)

Publication as pdf:

<http://www.umweltbundesamt.de/publikationen>

ISSN 1862-4804

Dessau-Roßlau, August 2022

The responsibility for the content of this publication lies with the author(s).

Table of content

List of figures.....	6
List of tables.....	7
Abbreviations.....	8
1 Background.....	9
2 An overview of ammonia research.....	11
2.1 Research in the early years.....	11
2.2 Research after 2009.....	15
2.2.1 Gradient Studies.....	18
2.2.2 Fumigation Studies.....	19
2.3 Ammonia monitoring.....	20
3 Discussion points at the workshop.....	32
4 Workshop presentations and discussion.....	33
4.1 Workshop Day 1, 28 March 2022.....	36
4.1.1 15 years on: Rationale and reflection on the 2006 Edinburgh Ammonia Workshop in the light of emerging evidence.....	36
4.1.2 Literature review on the effects of ammonia.....	47
4.1.3 Modelling ammonia concentrations, trends and scenarios in Europe.....	48
4.1.4 Measuring ammonia in the Netherlands.....	49
4.1.5 Continuous monitoring of ammonia concentrations with Cavity Ring-Down Spectroscopy.....	52
4.1.6 Ammonia monitoring at long-term and temporary sites in Flanders.....	53
4.1.7 Monitoring atmospheric ammonia on Natura2000 sites in the Republic of Ireland.....	54
4.1.8 National ammonia monitoring network in the UK: its current use and future potential.....	58
4.1.9 Ammonia concentrations measurements in Germany – an overview.....	63
4.1.10 Metrological aspects to support environmental ammonia monitoring.....	67
4.1.11 Long-term NH ₃ measurements at the TROPOS research station Melpitz using a MARGA.....	70
4.1.12 NADP’s ammonia monitoring network (Amon): Long-term measurements to support deposition and critical load assessments in the United States.....	72
4.2 Workshop Day 2, 29 March 2022.....	73
4.2.1 Relationships between NH ₃ in urban air and the diversity of epiphytic macrolichens.....	73
4.2.2 Results from an ammonia fumigation and a field study using endangered nitrogen-sensitive plant species.....	75

4.2.3	Towards understanding shorter-term critical levels with monthly resolution measurements	78
4.2.4	Biomonitoring of ammonia with the epiphytic lichen <i>Hypogymnia physodes</i>	81
4.2.5	How can we assess the impact of project contributions of ammonia on Natura2000 sites?	87
4.2.6	Development of a trophic assessment and management system for habitat types (Directive 92/43/EEC) considering interrelations between critical levels and critical loads.....	90
4.2.7	Ammonia concentrations in Switzerland compared to critical levels and loads	93
4.3	Workshop Discussion	95
4.3.1	Discussion points	95
4.3.2	Key findings.....	98
5	Acknowledgements	101
6	List of registered participants.....	102

List of figures

Figure 1:	Citations of Cape et al. (2009) in the peer-reviewed literature, broken down after years (top) and countries of residence of the main author (below)	16
Figure 2:	Numbers of presentations per country	33
Figure 3:	Comparison of ammonia concentration data from long term monitoring with the Egham 1992 critical levels for ammonia, as well as an extrapolated potential critical level for averaging over 25 years.	37
Figure 4:	Possible mechanisms by which atmospheric NH ₃ pollution affects epiphytic lichens, including both positive (+) and negative (-) effects.....	38
Figure 5:	Relationship between the Lichen Indicator Score (LIS) and the Nitrogen Air Quality Index (NAQI).....	39
Figure 6:	Illustration of the challenges of dealing with spatial scale in nitrogen deposition and critical loads exceedances. The landscape case study is taken from.	41
Figure 7:	Simulated a) annual global NH ₃ emissions (Gg N yr ⁻¹) from poultry farming in 2010. b) Percentage of excreted nitrogen that volatilizes (PV, %) as NH ₃ from chicken housing in 2010. The resolution is 0.5° 0.5°.	42
Figure 8:	Spatial distribution of intensive high-resolution miniDOAS (left) and low-cost Gradko (right) measurements in the Netherlands.	50
Figure 9	Ecological indicators of ammonia pollution on peatland site in the Republic of Ireland.....	55
Figure 10	Distribution of UK National Ammonia Monitoring Network sites	59
Figure 11:	Percentage of UK land area with ammonia concentrations exceeding critical levels: for a) lichens and bryophytes, b) vascular plants.....	60
Figure 12:	a) Map of annual mean NH ₃ surface concentrations over Germany (LOTOS-EUROS simulation for 2015; source: PINETI-3 project of the German Env. Agency). b) Annual box plot statistics of NH ₃ <i>in-situ</i> measurements from 2010-2019 for regional (upper panel) and remote (lower panel) locations.....	64
Figure 13:	Calibration strategies using gaseous reference standards (left) vs. an instrumental optical gas standard (right).	67
Figure 14:	Time series of the hourly ammonia concentration measured by a MAGRA over 10 years (a), the corresponding monthly average (b), and the seasonal diurnal patterns (c).....	70
Figure 15:	Shoot dry weight (g) of <i>E. cicutarium</i> , <i>B. hordeaceus</i> and <i>I. amara</i>	76
Figure 16:	Average annual NH ₃ concentrations at four designated nature conservation sites in Northern Ireland (top) and vegetation N	

	content (foliar ammonium) at four sites in Northern Ireland (bottom).....	79
Figure 17:	NH ₃ concentrations (4 – 6 monthly average) in Sandkrug (3 heights: 2m, 1.5m, 0.2m) and in Oldenburg (2m).....	82
Figure 18:	Exposure boards with <i>Hypogymnia physodes</i> in heights between 0.2 m and 2 m in Sandkrug.....	83
Figure 19:	Development of injury of <i>H. physodes</i> in Sandkrug within an exposure period of 2 years at different mean NH ₃ concentrations (confidence interval, medium, average, min. and max. value of injury in %).....	84
Figure 20:	Development of injury (%) of <i>H. physodes</i> within 15 months in Oldenburg (confidence interval, medium, average, min. and max. value) at mean NH ₃ concentrations of 2.5 µg m ⁻³	84
Figure 21:	Thalli of <i>Hypogymnia physodes</i> (left) and <i>Parmelia sulcata</i> (right) before (above) and after an exposure period of 15 months in Oldenburg.....	85
Figure 22:	Assessment scheme of nitrogen deposition impacts within German appropriate assessments under the Habitats Directive.....	87
Figure 23:	Ratio of road traffic contributions of ammonia- (yellow) and NO _x -depositions (green) at different distances along a road (forest).....	88
Figure 24:	Ammonia concentration and nitrogen deposition in Baden-Württemberg, average for the years 2012-2016.....	91

List of tables

Table 1:	Overview of ammonia fumigation experiments in closed and open chambers.....	13
Table 2:	Overview of ammonia monitoring programs.....	22
Table 3:	Summary of atmospheric ammonia (NH ₃) monitoring and modelled nitrogen (N) deposition across 12 Natura 2000 sites during 2017-2018 in the Republic of Ireland.....	54
Table 4:	The distribution of ammonia network sites by UK country (source: UKCEH).....	59
Table 5:	Considerations for an NH ₃ regulation in the EU AAQD.....	65
Table 6:	Results of the field comparison of the optical gas standard (OGS) and a mobile reference gas generator (ReGaS) (Twigg et al. 2022).....	68

Abbreviations

AAQD	Ambient Air Quality Directive
BImSchV	Bundesimmissionsschutzverordnung
CCE	Coordination Centre for Effects
CEH	Centre for Ecology and Hydrology (UK, several units)
CEN	Comité Européen de Normalisation
CLe	Critical Level for Air Pollutant Concentration, $\mu\text{g m}^{-3}$
CLo	Critical Load for Deposition, $\text{kg ha}^{-1} \text{yr}^{-1}$
CLRTAP	UNECE Convention on Long-Range Transboundary Air Pollution
CRDS	Cavity Ring-Down analyzers
DEFRA	Department for Environment, Food & Rural Affairs
DOAS	Differential Optical Absorption Spectroscopy
EMEP	The co-operative programme for monitoring and evaluation of the long-range transmission of air pollutants in Europe, European Monitoring and Evaluation Programme
IED	Industrial Emissions Directive
INI	International Nitrogen Initiative
INRA	Institut National de la Recherche Agronomique (France, several units)
ICP	International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects e.g. on Forests
IPCC	Intergovernmental Panel on Climate Change
IPPC	Integrated Pollution Prevention and Control
LANUV	Landesamt für Natur, Umwelt und Verbraucherschutz (LANUV) Nordrhein-Westfalen
LfU	Bayerisches Landesamt für Umwelt
LRTAP	UNECE Convention on Long-range Transboundary Air Pollution
LTER	Long-Term Ecological Research Network
LUBW	Landesanstalt für Umwelt Baden-Württemberg
MAK	Maximale Arbeitsplatz Konzentration (permissible workplace concentration)
NEC	National Emission Reduction Commitments Directive 2016/2284/EU
NERC	Natural Environment Research Council
OGS	Optical gas standard
PM	Particulate Matter PM_{10} , $\text{PM}_{2.5}$
PRTR	Pollutant Release and Transfer Register of the EU
QC-LAS	Quantum Cascade Laser Absorption Spectroscopy
ReGaS	Reference gas standard
RIVM	Rijksinstituut voor Volksgezondheid en Milieu (NL, Bilthoven)
TA Luft	Technische Anleitung zur Reinhaltung der Luft
UBA	Umweltbundesamt (German Environment Agency)
UNECE	United Nations Economic Commission for Europe
VDI	Verein Deutscher Ingenieure (The Association of German Engineers)
VMM	Vlaamse Milieumaatschappij (Flanders Environment Agency)

1 Background

Air pollution from ammonia (NH_3) is relevant for human health and ecosystems. The gas is easily converted into NH_4^+ , which can act as a plant nutrient, either by assimilation via the plant shoot (after entering the stomata) or after deposition to the ground and the subsequent uptake by the roots (Sutton et al. 1993; Fangmeier et al. 1994). Together with the oxidized forms of nitrogen, it is significantly contributing to the widespread eutrophication and acidification of ecosystems. As a result, the biodiversity of ecosystems is threatened at the expense of slow-growing nitrogen-sensitive, often rare and protected species (Bobbink et al. 2010).

Approximately 95 % of the ammonia (NH_3) emissions stem from agriculture, especially from livestock farming and the application of slurry or mineral fertilizers (Umweltbundesamt 2021). The odour threshold of ammonia is 5 ppm (3.8 mg m^{-3}) and the permissible workplace concentration (MAK) is 20 ppm (15 mg m^{-3}). Inside stables and after slurry application in the field, peak concentrations can be above 1 ppm ($760 \text{ } \mu\text{g m}^{-3}$), whereas own measurements have shown that outdoors and close to stables, mean levels will still be around $50 \text{ } \mu\text{g m}^{-3}$. Slightly elevated concentrations of the gas in the range of $10 \text{ } \mu\text{g m}^{-3}$ can also be detected near waste-water treatment plants. Major roads can act as line sources, but concentrations drop away from the edge of the road (Cape et al. 2004; Bell et al. 2016). In remote areas, ammonia concentrations are often below $1 \text{ } \mu\text{g m}^{-3}$ (Umweltbundesamt 2018), while the presence and excrements of animals (urea and uretic acid) and the degradation of plants will elevate levels locally.

Due to its alkaline character, ammonia is rapidly reacting with acids (e.g. sulphuric and nitric acid) creating ammonium salts like $(\text{NH}_4)_2\text{SO}_4$ and NH_4NO_3 , which possess long atmospheric lifetimes as secondary aerosols (PM 2.5) (Fowler et al. 2009). However, the strong reduction of the acidic air pollutants in the last decades has led to more alkaline atmospheric conditions and an increase in the lifetime of ammonia due to the reduced availability of H_2SO_4 to react with NH_3 (Sutton et al. 2020).

To protect European ecosystems from eutrophication, ammonia emissions will have to be reduced significantly in the EU according to the national emission ceilings laid down by the updated NEC Directive 2016/2284/EU and the 2020/2030 national emission reduction commitments. Most countries are currently developing and applying methods to assess nitrogen effects on ecosystems and vegetation with critical loads, in contrast critical levels are only used sporadically in the UK, Denmark and Germany within the framework of licensing new installations (Reinds et al. 2019).

Critical levels (CLE) describe the concentration of a pollutant above which direct adverse effects on receptors such as individual plants or natural ecosystems may occur. CLe for ammonia had been last discussed in the framework of the United Nations Economic Commission for Europe (UNECE) Convention on Long-range Transboundary Air Pollution (CLRTAP) at a Workshop on Atmospheric Ammonia (UNECE 2007). New experimental evidence was provided including from experiments at Whim Bog in Scotland (Cape et al. 2009) suggesting that the long-term critical level of ammonia needed to be reduced to $3 \text{ } \mu\text{g m}^{-3}$ for higher and $1 \text{ } \mu\text{g m}^{-3}$ for lower plants (i.e. lichens and mosses), respectively. The CLRTAP accepted these recommendations replacing existing the previous CLe of $8 \text{ } \mu\text{g m}^{-3}$ (Ashmore and Wilson 1994; UNECE 2007).

More than ten years after the recommendation of updated CLRTAP critical levels for ammonia in the Mapping Manual, new findings on the effects of ammonia on vegetation have been discussed on a workshop prepared by the CCE and the German Environment Agency (UBA) as a lead. The workshop was held at Dessau on March 28/29 March 2022 and offered the opportunity to exchange information on national or regional programs which have been set up for the monitoring of ammonia in sensitive habitats. Scientists dealing with research on effects of ammonia on vegetation

and ecosystems and those involved in the monitoring of ammonia in the environment were also asked to present their recent research.

The workshop was part of the current work plan for the year 2022-2023 of CLRTAP (1.1.1.22) and was organized in the framework of a R&D Project financed by the German Environment Agency (UBA). Due to the corona restrictions, only 15 participants were able to attend in person but more than 100 people were following the live stream via a WebEx meeting platform. Participants represented several European countries and the USA. In total, 19 talks were given from presenters from nine countries.

In the present workshop proceedings, we include a review of the latest findings and give summaries of the presentations given by the workshop participants.

2 An overview of ammonia research

Jürgen Franzaring¹, Matthew Jones^{2*}, Julia Köstler¹ and Alexander Moravek^{3*}

¹ University of Hohenheim, Institute of Landscape and Plant Ecology, Stuttgart, Germany

² UK Centre for Ecology & Hydrology, Edinburgh, United Kingdom, * co-authored chapter 2.2

³ German Environment Agency, Dessau, Germany, * co-authored chapter 2.3

In order to evaluate the validity of current critical levels and to prepare our own fumigation study (see Köstler & Franzaring, chapter 4.2.2), we screened the literature on ammonia effects from the last twelve years, focussing primarily on original publications which cited Cape et al. (2009). Nevertheless, we also reviewed important publications prior to 2009 in present background paper including fumigation studies from the 1970-1990s, that are helpful to better differentiate between phytotoxic vs. chronic effects and to address key points that need to be respected in fumigation experiments. Besides the peer-reviewed literature, we also included various reports from studies that monitored ammonia in regional and national projects.

2.1 Research in the early years

Experimental exposure of plants to ammonia in the very early years was mainly based on the use of pure (technical) NH₃ supplied from gas flasks to the air inlet of small closed fumigation chambers or continuously-stirred tank reactors (CSTR). Very high concentrations were applied initially to examine whether the growth of crop plants, e.g. sunflowers, could be stimulated in growth by nitrogen containing gases (Faller 1972; Ewert 1979; Temple et al. 1979; Farquhar et al. 1980). In these short-term experiments with young plants, phytotoxic responses (foliar injury and growth reductions) were observed only at concentrations above 800 µg m⁻³, depending on the plant species and the soil nitrogen supply. It had also been suggested early that fertilized crops and older plants do not take up ammonia any more at concentrations below 4 µg m⁻³, but release the gas during senescence (Cowling and Lockyer 1981).

Unfortunately, many of the early and more recent studies did not mention sufficient details how the ammonia fumigation was controlled, whether constant levels were achieved and how the realized concentrations were determined. Optimally, the relationships between the inlet and the outlet NH₃ concentrations would have to be monitored constantly (Jones et al. 2007). On the other hand, it will also be necessary to give a better account of how the test plants (receptors) were grown and what nutrient supply they received prior and during the fumigation. An overview of published ammonia fumigation experiments prior to 2009 is given in Table 1.

Dry deposition, bi-directional fluxes and compensation points of ammonia have been addressed early by Scottish, Dutch and Danish scientists (e.g. Sutton et al. 1993; Duyzer 1994 and Husted and Schjørring 1996) and have been parameterised in air-vegetation models later (Loubet et al. 2001; Nemitz et al. 2004). The fluxes are driven by changes to the equilibrium between the ammonia concentrations in the air and in the sub-stomatal cavity of plant leaves and depend on the air/leaf temperature, stomatal and cuticular resistances, seasonal and diurnal changes in stomatal opening, plant water status, ammonium (NH₄⁺) concentration (nutrient status) and pH of the apoplastic solution.

Farquhar et al. (1980) and van Hove et al. (1987) obtained compensation points of around 5 µg m⁻³ in crop plants, but suggest that in oligotrophic vegetation, e.g. peatlands, heath and dunes, the uptake of ammonia may occur at lower concentrations (e.g. Jones 2007b showed an average compensation point of >0.4 µg m⁻³ for moorlands at background atmospheric concentrations). This is principally confirming that low concentrations in the range of the suggested critical levels can be

“sensed” by vegetation in pristine areas, but as mentioned above, ammonia concentrations in agricultural and densely populated regions will only rarely be that low. During the lifetime and with the biological activity of plants during seasons, fluxes of ammonia will change. While young N-demanding plants are often a sink of ammonia in spring, old senescing plants may be a source and release the gas in the autumn. In addition, ammonia can be transported from a highly fertilized crop canopy to an adjacent natural vegetation with a lower compensation point and vegetation on acidic soil substrates will probably be more prone to the deposition of ammonia.

Following the initial discussion of critical levels for ammonia at the UNECE Workshop in Egham 1992 (Ashmore and Wilson 1994), European research networks had been established in the 1990s to monitor atmospheric processes that lead to the deposition of oxidized and reduced nitrogen species. It became clear that ammonia has a rather short life-time (Asman et al. 1998) reaching acute concentrations only close to emission sources. Further, chronic responses in sensitive wild plants will be driven by the joint deposition of various N-containing compounds mainly in the oligotrophic ecosystems. In order to account for the differences between short-term atmospheric concentrations and the cumulative annual inputs, the concept of critical loads had been introduced.

Ground-breaking research on the adverse effects and dose-response functions of ammonia on the natural (e.g. nitrogen-poor heathland) vegetation came from the Netherlands and served as the basis for the UNECE critical level of $8 \mu\text{g m}^{-3}$ (van der Eerden et al. 1982; Dueck 1990; van der Eerden et al. 1991; Dueck and Elderson 1992). Research of van der Eerden (1982) was based on the use of diluted exhaust from stables in closed or open top chambers (OTC), while later studies in Wageningen (Dueck and van der Eerden) involved technical ammonia supplied via mass flow controllers. Concentrations were in the range of 50 to $100 \mu\text{g m}^{-3}$ and were determined with NO_x converters/monitors. While there were several OTC experiments in the UK addressing NH₃ and wet deposition of N in the 1990-2000s, the only comparable research project performed in Germany dealt with an ammonia fumigation of three grassland species grown in monocultures and mixtures. The experiments were performed in OTCs with diluted technical ammonia using set concentrations of 20 and $50 \mu\text{g m}^{-3}$ (Jäger et al. 1998). However, the realized concentrations could not be documented due to failures of the NO_x converters/monitors. Summaries of the pioneering research of ammonia effects on plants can be found in the reviews of Adaros and Dämmgen (1994), Fangmeier et al. (1994) and Krupa (2003).

Based on the above-mentioned studies, we suggest that ammonia concentrations were probably too high in the chamber-based experiments from 1970 to ca 2000, restricting the validity of results with respect to the old and new critical levels. Expert judgements and extrapolations from high to low concentrations were used and long-term responses were predicted from short term experiments. None of the mentioned fumigation experiments operated in the range of existing environmental concentrations mainly due to technical reasons. When using pure ammonia without a time and concentration-controlled feedback system, peaks and accidental overdosing may occur that will lead to short-term burdens on plants eventually triggering acute effects. In fact, this will likely also happen in natural vegetation that is exposed to the fumes and plumes from adjacent slurry spreading or livestock housing.

Early Ammonia Research in growth chambers

- Focus on high concentrations and crop species, mainly in the Netherlands and the UK
- Mainly short-term experiments
- Pure (technical) NH₃ supply

Table 1: Overview of ammonia fumigation experiments in closed and open chambers

Source	Fumigation type	Supply of ammonia	Set range in $\mu\text{g m}^{-3}$	Method of ammonia measurement	Data on realized concentrations	Duration	Plant species
Faller (1972)	Glass chambers, 0.5 m ³	diluted NH ₃ , daylight hours	0 - 1600	none, calculated from flow	No	3 weeks	<i>Helianthus annuus</i>
Hutchinson et al. (1972)	Plexiglass chambers, 0.3 m high, 0.15 m sides	diluted NH ₃ ,	24-44	trapped in KCl solutions	24-44	24 hrs	<i>Glycine max</i> , <i>Zea mays</i> , <i>Helianthus annuus</i>
Ewert (1979)	2 chambers of 10 m ³	diluted NH ₃ , 24 hrs day ⁻¹	700 - 21000	air sampling, Nessler's reagent	No	1-500 hrs	48 woody species, 10 crops
Farquhar et al. (1980)	Glass chamber, 0.2 m ³	diluted NH ₃ , 24 hrs day ⁻¹	0 - 50 nbar	air sampling, indophenol method	No	20 minutes	<i>Phaseolus vulgaris</i> L. and other
Cowling & Lockyer (1981)	Exposure chambers	diluted NH ₃ , 16 hrs day ⁻¹	0, 148 and 550	air sampling, Nessler's reagent	No	40 days	<i>Lolium perenne</i> L.
van der Eerden (1982)	Open Top Chamber, closed chambers	Exhausts from poultry and pig manure, 24 hrs day ⁻¹	600 - 4500	air sampling, Nessler's reagent	No	3 - 60 days	grasses, trees, <i>Brassica</i> , <i>Lycopersicon</i>
Lockyer & Whitehead (1986)	Exposure chambers	diluted NH ₃ , 16 hrs day ⁻¹	0, 100 and 500	air sampling, indophenol method	16, 118, 520	40 days	<i>Lolium multiflorum</i> Lam.
van Hove et al. (1987)	Leaf Chamber	diluted NH ₃ , short term	0 - 400	Chemiluminescence after converting NH ₃ to NO	No	some days	<i>Phaseolus vulgaris</i> L.
Whitehead & Lockyer (1987)	Exposure chambers	diluted NH ₃	0 - 700	air sampling, indophenol method	14 - 709	33 days	<i>Lolium multiflorum</i> Lam.
Dueck (1990)	Open Top Chambers	diluted NH ₃ , 24 hrs day ⁻¹	0, 53 and 105	Chemiluminescence after converting NH ₃ to NO	No	9 months	<i>Calluna vulgaris</i>
Dueck & Elderson (1992)	Open Top Chambers	diluted NH ₃ , 24 hrs day ⁻¹	0, 50	Chemiluminescence after converting NH ₃ to NO	No	9 months	<i>Arnica montana</i> , <i>Viola canina</i> , <i>Agrostis capillaris</i>
Husted & Schjørring (1996)	Glass cuvette, 0.45 m ³	diluted NH ₃ , 24 hrs day ⁻¹	0 - 25 nmol mol ⁻¹	Chemiluminescence after converting NH ₃ to NO	No	twice a day, 7 days	<i>Brassica napus</i> L.
Jäger et al. (1998)	Open Top Chambers, 3.15 m diameter	diluted NH ₃ , 24 hrs day ⁻¹	20 and 50	Chemiluminescence after converting NH ₃ to NO	No	3 years	Monocultures and mixtures of <i>Bromus erectus</i> Huds, <i>Brachypodium pinnatum</i> P. Beauv. and <i>Arrhenatherum elatius</i> P. Beauv.
Geßler et al. (2002)	Borosilicate twig chambers	diluted NH ₃ , 24 hrs day ⁻¹	100	continuous wet flow denuder AMANDA	No	several hours	<i>Picea abies</i>
Adriral et al. (2006)	Exposure chambers	diluted NH ₃ , 24 hrs day ⁻¹	2800 - 5000	Photoacoustic NH ₃ detector	No	12 weeks	<i>Juniperus virginiana</i> , <i>Gleditsia triacanthos</i> , <i>Phalaris arundinacea</i> , <i>Populus spec.</i> <i>Salix spec.</i>

Table 1 continued

Source	Fumigation type	Supply of ammonia	Set range in $\mu\text{g m}^{-3}$	Method of ammonia measurement	Data on realized concentrations	Duration	Plant species
Jones et al. (2007a)	Flux chamber, 1.84 m ³ , placed in an OTC at Bush Estate near Edinburgh, Scotland	1% aqueous NH ₃ solution at different rates and concentrations	1-100	AMANDA system (Ammonia Measurement by ANnular Denuder sampling with on line Analyser system)	daily resolution	6 hours each trial over 3 to 6 weeks	mixed turfs made up of <i>Calluna vulgaris</i> (L.) Hull, <i>Eriophorum vaginatum</i> L., <i>Sphagnum capillifolium</i> (Ehrh.) Hedw. and <i>Sphagnum papillosum</i> Linb.
Jones et al. (2007b)	Flux chamber, 1.84 m ³ , placed inside of an OTC	1% aqueous NH ₃ solution at different rates and concentrations	1-140	continuous wet flow denuder AMANDA, measurements at inlet and outlet	daily resolution	6 hours each trial over 3 to 6 weeks, stomatal and non-stomatal species	<i>Deschampsia cespitosa</i> (L.) Beauv., <i>Calluna vulgaris</i> (L.) Hull, <i>Eriophorum vaginatum</i> L., <i>Cladonia</i> spp., <i>Sphagnum</i> spp., and <i>Pleurozium schreberi</i> (Brid.)
Jones et al. (2008)	Flux chamber, 1.84 m ³	release of ¹⁵ N labelled NH ₃ from (¹⁵ NH ₄) ₂ SO ₄ with NaOH	16 - 20	continuous wet flow denuder AMANDA	daily resolution	15 days	<i>Calluna vulgaris</i>
Franzaring et al. (2012)	Mini greenhouses	diluted NH ₃ , 24 hrs day ⁻¹	0 - 200	Passive samplers	2.2 -195	88 days	<i>Brassica napus</i> L.
Ilogu Chibuzo (2012)	Exposure chambers	controlled evaporation of NH ₄ Cl solutions	0 - 84	Passive samplers	3, 25, 84	30 days	<i>Echinochloa crus-gallii</i> and <i>Lolium multiflorum</i>
Jones et al. (2013), presenting results from PhD Thesis of Jonathon Foot from 1998	Open Top Chambers	uncontrolled evaporation of NH ₄ OH solutions containing 0-10 % of ammonia	0 - 35	Passive samplers	Yes	28 weeks	Mixtures of <i>Dactylis glomerata</i> , <i>Plantago lanceolata</i> , <i>Festuca rubra</i> , <i>Centaurea nigra</i> etc.

2.2 Research after 2009

It must be highlighted that during the last 30 years many research activities on the fluxes, deposition and effects of ammonia came from Scotland, namely David Fowler, Neil Cape and Mark Sutton, all scientists from the Centre for Ecology & Hydrology (CEH) Edinburgh. These authors have co-authored many studies and worked as PI's of several EU projects that dealt with the deposition of acidifying and eutrophying gases and particulates. We only cited a few of these studies in previous chapter, but acknowledge the impact they still have on current research.

In present sub-chapter we will take a closer look at research that deals with the effects of plants and the vegetation under more natural conditions. Whereas chamber-based experiments (see. 2.1) dealing with plant exposure to ammonia were mostly applying higher concentrations, more recent projects tried to investigate low-range ammonia effects in order to address potential responses in sensitive plant species and to be more representative for the range of critical levels.


A screening of newer literature dealing with the recommended CLe and contributing new data from the UK and other European countries will be presented in this subchapter. We present an overview of 103 peer-reviewed references (as of December 2020) that cited Cape et al. 2009 according to the databank Scopus®. Figure 1 (top) gives an overview on how many citations per year were recorded and how many of these did not stem from the Scottish research group that also authored the Cape et al. publication. After a decline in the numbers of citations until 2019, the topic “critical levels” peaked again recently and we can observe that 80 % of the citations were from groups not involved in writing the cited paper.

When looking at the country of residence of the main author and her affiliation (Figure 1 below), we can see that most publications which cited Cape et al. (2009) stemmed from the UK, China, Canada and the US, being followed by three Mediterranean countries mostly studying lichen diversity among ammonia gradients. Interestingly, in the Netherlands, Denmark and Germany, the research on effects of ammonia is not of major importance, although the countries have a very large livestock production sector and severe problems with eutrophication. It must be noted that until the late 1990s, most of the ground-breaking research on ammonia impacts had been performed in the Netherlands (the research group of Dr. Ludger van der Eerden at Plant Research International in Wageningen). The old CLe of $8 \mu\text{g m}^{-3}$ had originally been derived from fumigation studies from that institute and in those years, there were many ammonia-related research programs funded by the Dutch, UK and German governments. An even stronger absence of research was noted in France, a country with a strong livestock sector in the EU, where no peer-reviewed literature on the effects of ammonia was published after 2009.

When looking at the specific topics dealt with in the 103 articles, the study of the presumed ammonia sensitivity of lichens plays an overwhelming role. 27 articles focus on the mapping of nitrogen sensitive lichens with many published from Southern European countries, despite the limited intensive livestock. Another quarter of the citing articles dealt with the modelling of N deposition and 20 articles were presenting data from ammonia measurements using passive samplers. 16 studies investigated changes in the species composition. 11 studies reported about “bioindication” methods, addressing ecophysiological changes in lower and higher plants along ammonia gradients. Only a few of these studies used pre-cultivated higher plants, which were actively exposed in the field at locations, where ammonia was measured. In the recent paper of Ellis et al. (2022) on potential nitrogen risks to Himalayan forests, the results of 14 lichen biomonitoring studies were used to derive independent estimates for three nitrogen pollutant types. These were the critical level for ammonia concentration ($\mu\text{g m}^{-3}$), the critical load for total nitrogen deposition ($\text{kg N ha}^{-1} \text{ yr}^{-1}$) and the total throughfall nitrogen deposition. Based on their calculations for diverse responses on lichens (e.g. total species richness, sensitive species richness, ‘forage lichen’

abundance and cyanolichen abundance), the critical level for ammonia would be $1.44 \mu\text{g m}^{-3}$ as a grand mean and the critical load (wet and dry, reduced and oxidized) would average to $8.26 \text{ kg N ha}^{-1} \text{ yr}^{-1}$.

Figure 1: Citations of Cape et al. (2009) in the peer-reviewed literature, broken down after years (top) and countries of residence of the main author (below)



Source: Based on a literature search in the SCOPUS database © Elsevier B.V

Besides the agricultural sources, the NO_x emissions from industry, traffic and wastewater treatment sectors are also responsible for the losses in biodiversity of certain cryptogams and lichens. Sutton et al. (2020) have recently suggested that the successful reduction of SO₂ pollution in Europe and elsewhere has led to a general alkalinisation of rain and air that will be harmful to naturally acidophytic species. On the other hand, in the 1990s and early 2000s the elimination of acidification often favoured the return of rare basiophytic lichens. The impacts of the recent and ongoing changes in air quality have also been investigated in Sweden and showed a very slow recolonization of sensitive species due to existing dispersal limits (Weldon and Grandin 2021).

Only in the 1990s, research on the phytotoxicity of air pollutants moved to field-based fumigation systems to study the effects of ammonia on pre-cultivated plants or established natural vegetation. Such systems should operate under the normal ambient climates and should not create concentration peaks. Examples are the FACE (free air carbon dioxide enrichment) experiments and the use of open top chambers (OTC) e.g. to study adverse effects of ozone.

The only long-term field ammonia fumigation system is at the Scottish Whim Bog Field Facility (Southwest of Edinburgh) and has been in operation since 2002. The site simulates a point source release (e.g. chicken farm) producing a concentration gradient across an ombrotrophic bog, ranging from an annual average of around $80 \mu\text{g m}^{-3}$ (at 6 m) to ambient ($<1 \mu\text{g m}^{-3}$). The site represents a release of approximately 5.000 chickens in a clean shed system (SCAIL model estimate) and is similar to annual concentrations reported near comparable farms (e.g. Pitcairn 1998). All major species have declined (including *Cladonia*, *Sphagnum*, *Calluna*, *Hypnum*) except the now dominant sedge *Eriophorum vaginatum*, across the large majority of the concentration ranges (van Dijk et al. 2017; Levy et al. 2019). Evidence suggests long term effects with impacts responding to changes occurring over several years (Sutton et al. 2020). Severe impacts of lichen bleaching recorded near the source may be caused by peak toxicity events, but impacts along the transect are likely caused by other mechanisms such as algal/fungal imbalances.

More recently in 2021, CEH has established a new field ammonia fumigation system at Glencorse Field Facility (Southwest of Edinburgh). This site releases ammonia at a similar level to Whim, but assesses the impact of ammonia within a woodland environment (pers. comm. Matthew Jones). The site has 20 m long fumigation pipes at three heights to ensure ammonia is relatively well mixed in the trunk region of the woodland. Ammonia is only released when wind directions and speed are adequate to ensure maximum concentrations are representative of concentrations expected downwind of intensive agriculture. Significant pre-testing of the system was conducted using a Picarro gas concentration NH_3 analyser in a multi-height inlet system, multi-height profiles of ALPHA samplers, a transect of ALPHAs and ALPHAs placed throughout the woodland to allow the researchers to establish the plume shape and concentrations under different meteorological conditions. The site also has a 15 m high meteorological mast to establish wind, temperature, relative humidity and leaf wetness profiles. The site is being used as a test bed to improve – model estimates of NH_3 deposition to woodlands, NH_3 impacts on forest ecosystems, NH_3 impacts on GHG emissions, and the work will include assessment of NH_3 impacts on lichen transplants in relation to critical levels. A third site has also now been established (March 2022) in Sri Lanka following the same intensive pre-measurements as at the Glencorse experiment, which will for the first time assess nitrogen impacts in South Asian ecosystems, and will be compared to a large-scale assessment of N impacts in the Himalayan Forests in South Asia.

A new generation of experiments mimicking the dry gaseous deposition of ammonia in established ecosystems is currently on the way also in China (Pan et al. 2020) and will be recognized under the name Free-Air NH_3 Enrichment (FANE).

Critical Levels Research

- Until 2020, Cape et al (2009) had been cited over 100 times
- Most citations stem from UK, China, Canada and the US
- Relevant number of publications in several Mediterranean countries
- Low number of publications from the Netherlands, Denmark, Germany and France, although the countries have a large livestock sector and eutrophication problems are common
- Citing studies deal with lichen mapping, modelling N deposition, ammonia measuring, changes in species composition and the investigation of ecophysiological changes in lower species
- Critical loads research is more prominent than the study of ammonia or nitrogen dioxide

2.2.1 Gradient Studies

Most field experiments dealing with the effect of ammonia used passive approaches working with the reaction of established vegetation on ammonia input. Sheppard et al. (2011) described a dramatic loss of higher and lower plants at Whim Bog after three years of relatively modest NH₃ deposition, with impacts after 14 years of treatment being observed down to low NH₃ concentrations (Levy et al. 2019). Several lichen mapping studies also showed effects of ammonia with regard to lichen diversity along natural gradients. The number of oligotrophic lichen species decreased significantly with increasing atmospheric ammonia concentrations whereas nitrophytic species increased and/or replaced some species (Pinho et al. 2011; Pinho et al. 2012; Pinho et al. 2014). Pinho et al. suggested CL_e of ammonia for Mediterranean evergreen woodlands to be 1.9 µg m⁻³ (2012) and two years later they suggested a lower level of 0.69 µg m⁻³. Aguilhaume et al. (2017) suggested a higher CL_e of 2.6 µg m⁻³ for a Mediterranean Holm-Oak Forest in Spain, but argue that the site has already been polluted by NH₃ in the past. Another study from Spain monitored NH₃ in peri-urban forests (*Quercus ilex*) and found no CL_e exceedance in any of the three sites (García-Gómez et al. 2016). It was furthermore shown, that below-canopy concentrations of NH₃ were around 40% lower than in open-field plots, indicating a high sink capacity of holm oak forests.

Mayer et al. (2013) showed that lichen diversity in Austria has been declining in a remote area (in the European Alps) and suggest eutrophication due to ammonium in precipitation and fog as an important factor. Experiments from Canada revealed that most lichen species studied were affected above 1.5 µg m⁻³ (Watmough et al. 2014). Izquieta-Rojano et al. (2018) describe effects of an ammonia gradient (distance to a swine livestock facility) on mosses (especially tissue N content) in an oak woodland with an estimated site-specific CL_e of 3.5 µg m⁻³. The results imply that mosses might stand higher concentrations of ammonia than lichens, but it must be noted that there is less information on mosses compared to lichens at the moment. Other passive approaches like the use of oak leaf characteristics as a biomonitoring tool. (Wuytack et al. 2013) revealed no differences and therefore seem not to be suitable for ammonia monitoring programs.

Active (biomonitoring) studies are mainly addressing the effects of ammonia from agricultural sites on lower plant species, i.e. lichen and/or moss diversity and vitality. Sensitive moss (Freiberger et al. 2021) or lichen species like *Evernia prunastri* (Paoli et al. 2010; Paoli et al. 2014; Munzi et al. 2014) or *Xanthoria parietina* and *Flavoparmelia caperata* (Fрати et al. 2007) had been transplanted from non-polluted sites (sites with low ammonia background concentrations) to natural ammonia gradients and the vitality e.g. in form of chlorophyll fluorescence had been determined. Munzi et al. (2014) found that ammonia concentrations above 3 µg m⁻³ would abruptly decrease plant fitness and frequency but the authors approve the same NH₃ CL_e (1 µg m⁻³) as proposed by Cape et al (2009) because of possible cumulative effects after longer exposure than tested. Interestingly, a more recent study observed that the N-sensitive lichen *Cladonia portentosa* did not show any adverse effects anymore after ceasing the wet deposition of nitrogen (Munzi et al. 2020).

The effects of a higher availability of nitrogen along gradients do not only depend on the form of N and the wet/dry or gaseous/particulate deposition but are also affected by environmental gradients. Morillas et al. (2021) studied detrimental cumulative effects of solar radiation and increased nitrogen treatments on lichens via immersing samples into N solutions. Studies about ammonia effects are rarely including additional climatic factors to date. Freiberger et al. (2021) recently suggested that active approaches using uncontaminated moss transplants (*Pleurozium schreberi*) and accumulated N-levels show a much stronger relationship to the ammonia concentrations than moss or lichen mapping (passive approaches).

Gradient studies in the vicinity of specific emitters making use of higher plants are often focussing on grass species as bioindicators since their upright growth habit and the presence of numerous stalks enable the effective scavenging and accumulation of air pollutants. While Leith et al. (2009) used grasses in the lee of Scottish poultry farms (confirming CLe of 2-3 $\mu\text{g m}^{-3}$), Ilogu Chibuzo (2012) used grasses and N-demanding weeds in the lee of a pig stable in Southern Germany. In both studies, plant biomass increments and nitrogen concentrations proved to be sensitive indicators for the fertilizing effect of ammonia.

Gradient Studies

- Most field studies make use of passive approaches, i.e. investigate the reaction of established vegetation on ammonia concentrations in a natural gradient
- Many studies show a decrease of N-sensitive lichen species with increasing NH_3 concentrations
- Mosses seem to be more tolerant than lichens
- Most active study approaches with lichen transplants confirm low CLe
- Most recent studies addressing CLe for ammonia work on passive approaches with lichens in natural ammonia gradients, the majority suggesting CLe of 0.69-3.5 $\mu\text{g m}^{-3}$
- Only few studies suggest that lichens might stand higher NH_3 concentrations than expected
- Only very few studies address higher plants
- Active approaches confirm CLe of 1 $\mu\text{g m}^{-3}$ for lichens and 2-3 $\mu\text{g m}^{-3}$ for higher plants (e.g. grasses)

2.2.2 Fumigation Studies

Apart from the free-air exposure at Whim Bog, which uses the release of technical ammonia along a wind gradient, only two peer-reviewed papers on ammonia fumigation could be found that had been published after 2009. In our own experiment (Franzaring et al. 2012), we studied the effects of sub-acute concentrations of ammonia on the re-growth of winter oilseed rape. We could show that the crop is well able to benefit from the gaseous fertilizer at a concentration of 195 $\mu\text{g m}^{-3}$, but we did not test whether slightly raised or low concentrations of ammonia would be able to significantly interact with the physiology of the species.

The second study on ammonia fumigation was published by Jones et al. (2013). The authors investigated the effects of ammonia deposition to plants from Welsh sand dune ecosystems and exposed mesocosms of plant mixtures in OTCs. Shoot biomass and tissue nitrogen increased significantly with increasing concentrations of ammonia. A significant biomass increase of the plant mixture was already observed when increasing the ammonia concentrations from 0.4 to 4.3 $\mu\text{g m}^{-3}$, i.e. dose equivalents of 2 and 24 $\text{kg N ha}^{-1} \text{y}^{-1}$.

Fumigation Studies

- Apart from the Whim Bog Experiment, only two newer fumigation studies were found
- Both studies showed a biomass increase of the species tested under NH_3 fumigation

2.3 Ammonia monitoring

Since the 1990s, pioneering work on ammonia measurements in the ambient air has been done in the Dutch Landelijk Meetnet Luchtkwaliteit (LML) and in the UK National Ammonia Monitoring Network (NAMN). Various types of denuders and passive sampler devices are being applied to allow for a resolution ranging from days to weeks.

More recently, other devices have been made available for fast, continuous, real-time monitoring of ammonia. Examples are devices based on Differential Optical Absorption Spectroscopy (DOAS), Cavity Ring-Down Spectroscopy (CRDS) or Quantum Cascade Laser Absorption Spectroscopy (QCLAS). While those methods are well able to determine low background concentrations, they are typically not employed in monitoring networks due to their high cost and high demand for support. In the Netherlands and in Switzerland a few monitoring sites employ DOAS devices (Volten et al. 2012a; Volten et al. 2012b; Berkhout et al. 2017; Bell et al. 2017) and more CRDS instruments may be included in monitoring initiatives in the near future (see Pogány et al., Poulain and van Zwieten, in this issue).

A major challenge in measuring ammonia concentrations is its stickiness to surfaces of measurement devices. Therefore, there is a high need for more intensive metrological research focusing on the quality assurance, inter-comparability and validation of ammonia in the ppb range.

A first international initiative comparing different active and passive analysers was set up within the European MetNH₃ (Metrology for ammonia in ambient air) project. One of the work packages of the MetNH₃-Project dealt with the testing and calibrating of ammonia passive samplers. Samplers were tested in the UK's National Physical Laboratory's (NPL) controlled atmosphere test facility (CATFAC) and in the field at the above-mentioned Scottish Whim Bog Experimental site. The inter-comparisons between different sampler types have been presented by Braban et al. (2018) and Martin et al. (2019). Measurement errors of the tested passive samplers varied between 11 and 23 % at the reference concentration of 1 µg m⁻³, i.e. a concentration representative for the lower critical level. ALPHA passive samplers proved to be the best suited for the lower concentration ranges and have an exposure time of 28 days, whereas e.g. the Radiello® samplers can be used to investigate shorter periods.

Most recently, a comprehensive comparison of measurement techniques under field conditions has been presented by Twigg et al. (2022) and will be discussed by the research community.

Here we focus mainly on the results determined by passive samplers. These cost-effective devices have been developed for the determination of gaseous air pollutants in locations without power supply and are therefore the preferred method to monitor gas concentrations in remote areas. In order to assure their quality, devices should be certified according to CEN (European Committee for Standardization) standards EN 17346 (2020)¹ or comparable national standards for diffusive samplers. Two types are available, axial (badge and tube types) and radial (tube like) samplers. Latter types have higher sampling rates and can therefore be exposed over shorter times (Puchalski et al. 2011).

Table 2 gives an overview of studies that used passive samplers to study ammonia concentrations in different countries. Only papers and grey literature are listed that were published after 2009. In most of the studies, ammonia concentrations did not decrease over time contradicting the results from modelled concentrations (Wichink Kruit et al. 2017) and the information derived from emission inventories. As previously mentioned, the reasons for higher levels may be the reduced concentrations of NO_x and SO₂ and increased volatilization of the gas due to the higher ambient

¹ <https://www.en-standard.eu/csn-en-17346-ambient-air-standard-method-for-the-determination-of-the-concentration-of-ammonia-using-diffusive-samplers/>

temperatures. At the same time, higher concentrations may also stem from previously underestimated emissions, e.g. due to uncertainties in emission factors or activity data in agriculture, from waste water treatment facilities and biogas plants as well as from the increased use of diesel exhaust fluids (DEF, AdBlue). Higher vehicular emissions of NH_3 and ammonia slip have been addressed by Elser et al. (2018), Suarez-Bertoa and Astorga (2016) and Fenn et al. (2018).

In contrast to the pioneering, smaller European countries like the Netherlands (Lolkema et al. 2015; Noordijk et al. 2020), Belgium (VMM 2017) and Switzerland (Thöni et al. 2018; Seitler & Meier 2022) and the UK (Tang et al. 2018), countries with intensive agricultural systems like France, Italy and Germany do not operate national ammonia networks yet, so that regional data are scarce in these countries. We could only find one publication from France that reports on ammonia measurements in five agricultural and six forest regions (Fauvel et al. 2019). In Brittany, a region with intensive livestock farming, ammonia concentrations are on average in the range of $9 \mu\text{g m}^{-3}$, while inside forest areas in the south and in the east, concentrations of less than $1 \mu\text{g m}^{-3}$ are common. In Germany, there have been some efforts in various federal states to monitor ammonia with passive samplers, mainly badge type devices (e.g. Lohrengel et al. 2012; LfU 2019).

Belgium, the Netherlands and Ireland have established monitoring sites in a large number of Natura2000 (the EU Habitat Directive) areas (Table 2). In contrast, in Germany measurements are being performed on a voluntary basis and are not conducted with uniform objectives in eleven out of sixteen Federal States. Since protected landscapes are often nitrogen limited, the vegetation (e.g. bogs and heathlands) is an important receptor for potentially harmful ammonia. In the present workshop proceedings, results from some European countries and Northern America will be presented in greater detail (see the contributions of Staelens, Meier, Kelleghan, Gay & Puchalski, Moravek & Geupel, this issue).

Ammonia Measuring and Monitoring

- Ammonia monitoring programmes have been set up in both, N-America and Europe. Also, in China, NH_3 is being determined at many locations of the national N deposition networks.
- The intensity and technology used for ammonia monitoring differs between countries. The Netherlands and Switzerland are routinely operating a few sites with high-precision trace gas measurements, while in other countries DOAS and CRDS are applied in short-term research projects only.
- Many countries and studies use denuders and passive samplers, latter method representing a well-accepted, feasible low-cost technique to measure NH_3 .
- Real-time monitoring gas analysers are mainly used in temporary, small-scale initiatives. Their continuous development towards lower maintenance requirements and easier handling may make them more suitable for continuous monitoring of ambient levels of ammonia in the future.

Table 2: Overview of ammonia monitoring programs

List is based on knowledge of monitoring programs to authors at the time of publication and may be incomplete.

Country	Sampler type	Network name	N of sites ²	Period	Network Details / Results	Sources
BE	Radiello®, blue diffusion barriers, miniDOAS		100 Natura2000 sites, miniDOAS at 2 sites	2015-2016	Country mean: 3.3 µg m ⁻³ NH ₃ , in Western Flanders > 5 µg m ⁻³ NH ₃ , no trends over time.	VMM (2017)
CAN-ONT	Ogawa	NAPS	74	2006-2007	Even in remote sites in CAN, 50% of the sites would exceed the critical levels.	Yao & Zhang (2013; 2019)
CH	Radiello®		13	since 2000	No reductions over time, different levels in different regions. All values exceed the 1 µg level, most the 3 µg critical level.	Seitler et al. (2018); Seitler & Meier (2022)
DE -Depo	FERM (IVL)		15 forest sites (in 2020)	since 2001	ICP-Forests Level II locations, bi-weekly. No decline in concentrations in German Forests. Grand mean: 1.9 µg m ⁻³ .	Thünen Institute of Forest Ecosystems (pers. comm.)
DE-BW	FERM (IVL)		16 ³	since 2007	Selected urban and rural locations. Moderately high NH ₃ at traffic sites.	LUBW (https://pd.lubw.de/10334)
DE-BY	Ferm (IVL)	LÜB, DBS	15	since 2011	Selected locations, four-weekly. Downward trend in cities, in rural areas increasing trend since 2017. Yearly mean levels up to 3 µg m ⁻³ at rural sites, 7 µg m ⁻³ at intensive agricultural sites.	LfU (2019) and LfU (pers. comm.)
DE-MV			10	2010	No trend study possible.	https://www.lung.mv-regierung.de/umwelt/luft/archiv/jb_2019_small.pdf
DE-NRW	Ferm (IVL)		5	since 2001	Selected urban and rural locations. No trend.	Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen (pers. comm.)
DE-NI	Ferm (IVL)		19	since 2010	Selected rural locations. 0.7 to 13 µg m ⁻³ ammonia. Two sites in industrial settings.	https://www.umwelt.niedersachsen.de/download/183444/Jahresbericht_2021.pdf
DE-SH			10	2008	No trend study possible.	https://www.schleswig-holstein.de/DE/fachinhalte/Luftqualitaet/Downloads/Berichte/bericht_orient_messungen_NH3_SH_2011.pdf?blob=publicationFile&v=1

² If measurements are still ongoing, number represents the current number of sites with operating ammonia measurements

³ BW established 13 new sampling sites in 2021. They are not officially documented, therefore not reported here

Country	Sampler type	Network name	N of sites	Period	Results	Sources
DE-ST			7	since 2021/22	No trend study possible.	LUA (https://lau.sachsen-anhalt.de/luft-klima-laerm/lufthygienisches-ueberwachungssystem-luesa/)
DE-UBA	Denuder		7	2004	Network background sites of the German Environmental Agency	UBA (https://www.umweltbundesamt.de/das-uba/standorte-gebaeude#dessau)
F	Alpha, Delta		11	2009-2019	Selected agricultural and forest locations. High ammonia concentrations in Brittany.	Fauvel et al. (2019)
IRL	Alpha		12 Natura2000 sites	2017-2018	0.47–4.59 $\mu\text{g NH}_3 \text{ m}^{-3}$. Agriculture is the main source and ammonia is related to ambient temperatures.	Kelleghan et al. (2021)
LUX	Gradko		4	since 2019	Mean concentrations vary from 5.5 to 19.2 $\mu\text{g m}^{-3}$ at rural and agricultural sites.	Pers. comm. Trebs & Junk, Luxembourg Institute of Science and Technology (LIST 2022)
NL	Gradko, miniDOAS	MAN	Currently 310 measuring points at 105 sites, out of which 77 are Natura2000 sites, mini DOAS operated at 6 sites	since 2005	NH_3 concentrations significantly increased in easterly regions and went down in the west and in the south.	Lolkema et al. (2015); Noordijk et al. (2020); pers. comm. Jonkers and Bleeker
UK	Alpha, Delta	UK NAMN	74	since 1998	Reduction of 3% on national level, but ammonia increases in some regions.	Tang et al. (2018); United Kingdom Eutrophying & Acidifying Network (UKEAP), https://uk-air.defra.gov.uk/networks/
US	Radiello®, blue diffusion barriers	AMoN	50	since 2007	NH_3 concentrations have increased by 7% between 2008 and 2015 at 18 long-term sites.	Puchalski et al. (2011) and Yao & Zhang (2016; 2019)

References

- Adaros, G.; Dämmgen, U. (1994): Phytotoxische Wirkungen der aktuellen NH₃-Immissionen. Bundesforschungsanstalt für Landwirtschaft. Braunschweig 1994.
- Aguillaume, L.; Avila, A.; Pinho, P.; Matos, P.; Llop, E.; Branquinho, C. (2017): The Critical Levels of Atmospheric Ammonia in a Mediterranean Holm-Oak Forest in North-Eastern Spain. In *Water, Air, & Soil Pollution* 228 (3), p. 919. DOI: 10.1007/s11270-017-3286-8.
- Adrizal; Patterson, P. H.; Hulet, R. M.; Bates, R. M. (2006): Foliar Nitrogen Status and Growth of Plants Exposed to Atmospheric Ammonia (NH₃). Workshop on Agricultural Air Quality
- Ashmore, M. R.; Wilson, R. B. (1994): Critical levels of air pollutants for Europe. Background Papers prepared for the United Nations Economic Commission for Europe Workshop on Critical levels. Egham UK, 23-26 March 1992, 209 pp.: Air Quality Division, Department of the Environment, London.
- Asman, W. A. H.; Sutton, M. A.; Schjoerring, J. K. (1998): Ammonia: emission, atmospheric transport and deposition. In *New Phytologist* 139, pp. 27–48.
- Bell, M.; Flechard, C.; Fauvel, Y.; Häni, C.; Sintermann, J.; Jocher, M. et al. (2017): Ammonia emissions from a grazed field estimated by miniDOAS measurements and inverse dispersion modelling. In *Atmospheric Measurement Techniques* 10 (5), pp. 1875–1892. DOI: 10.5194/amt-10-1875-2017.
- Bell, M. W.; Tang, Y. S.; Dragosits, U.; Flechard, C. R.; Ward, P.; Braban, C. F. (2016): Ammonia emissions from an anaerobic digestion plant estimated using atmospheric measurements and dispersion modelling. In *Waste Management (New York, N.Y.)* 56, pp. 113–124. DOI: 10.1016/j.wasman.2016.06.002.
- Berkhout, A. J. C.; Swart, D. P. J.; Volten, H.; Gast, L. F. L.; Haaima, M.; Verboom, H. et al. (2017): Replacing the AMOR with the miniDOAS in the ammonia monitoring network in the Netherlands. In *Atmospheric Measurement Techniques* 10 (11), pp. 4099–4120. DOI: 10.5194/amt-10-4099-2017.
- Bobbink, R.; Hicks, K.; Galloway, J.; Spranger, T.; Alkemade, R.; Ashmore, M. et al. (2010): Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. In *Ecological applications: a publication of the Ecological Society of America* 20 (1), pp. 30–59. DOI: 10.1890/08-1140.1.
- Braban, C. F.; Bree, F. de; Crunaire, S.; Fröhlich, M.; Fromage-Mariette, A.; Goelen, E. et al. (2018): Literature review on the performance of diffusive samplers for the measurement of ammonia in ambient air and emissions to air. Edinburgh, Centre for Ecology & Hydrology, 85pp (CEH Project no. C05204, C05967, C04544, C05952, C06942) (Unpublished).
- Cape, J. N.; Tang, Y. S.; van Dijk, N.; Love, L.; Sutton, M. A.; Palmer, S. C. F. (2004): Concentrations of ammonia and nitrogen dioxide at roadside verges, and their contribution to nitrogen deposition. In *Environmental Pollution (Barking, Essex: 1987)* 132 (3), pp. 469–478. DOI: 10.1016/j.envpol.2004.05.009.

Cape, J. N.; van der Eerden, L. J.; Sheppard, L. J.; Leith, I. D.; Sutton, M. A. (2009): Evidence for changing the critical level for ammonia. In *Environmental Pollution* (Barking, Essex: 1987) 157 (3), pp. 1033–1037. DOI: 10.1016/j.envpol.2008.09.049.

Cowling, D. W.; Lockyer, D. R. (1981): Increased growth of ryegrass exposed to ammonia. In *Nature* 292 (5821), pp. 337–338. DOI: 10.1038/292337a0.

Dueck, Th. A. (1990): Effect of ammonia and sulphur dioxide on the survival and growth of *Calluna vulgaris* (L.) Hull Seedlings. In *Functional Ecology* 4 (109-116).

Dueck, Th. A.; Elderson, J. (1992): Influence of NH₃ and SO₂ on the growth and competitive ability of *Arnica montana* L. and *Viola canina* L. In *New Phytologist* 122, pp. 507–514. DOI: 10.1111/j.1469-8137.1992.tb00080.x.

Duyzer, J. (1994): Dry deposition of ammonia and ammonium aerosols over heathland. In *Journal of Geophysical Research* 99 (D9), pp. 18757–18763. DOI: 10.1029/94JD01210.

Ellis, C. J.; Steadman, C. E.; Vieno, M.; Sudipto, C.; Jones, M. R.; Negi, S. et al. (2022): Estimating nitrogen risk to Himalayan forests using thresholds for lichen bioindicators. In *Biological Conservation* 265, p. 109401. DOI: 10.1016/j.biocon.2021.109401.

Elser, M.; El-Haddad, I.; Maasikmets, M.; Bozzetti, C.; Wolf, R.; Ciarelli, G. et al. (2018): High contributions of vehicular emissions to ammonia in three European cities derived from mobile measurements. In *Atmospheric Environment* 175 (Suppl. 1), pp. 210–220. DOI: 10.1016/j.atmosenv.2017.11.030.

Ewert, E. (1979): Zur Phytotoxizität von Ammoniak. In *Hercynia* N. F. 16 (1), pp. 75–80.

Faller, N. (1972): Schwefeldioxid, Schwefelwasserstoff, nitrose Gase und Ammoniak als ausschließliche S- bzw. N-Quellen der höheren Pflanze. In *Journal of Plant Nutrition and Soil Science* 131 (2), pp. 120–130. DOI: 10.1002/jpln.19721310204.

Fangmeier, A.; Hadwiger-Fangmeier, A.; van der Eerden, L. J.; Jäger, H.-J. (1994): Effects of atmospheric ammonia on vegetation - A review. In *Environmental Pollution* 86 (1), pp. 43–82. DOI: 10.1016/0269-7491(94)90008-6.

Farquhar, G. D.; Firth, P. M.; Wetselaar, R.; Weir, B. (1980): On the Gaseous Exchange of Ammonia between Leaves and the Environment: Determination of the Ammonia Compensation Point. In *Plant Physiology* 66, pp. 710–714. DOI: 10.1104/pp.66.4.710.

Fauvel, Y.; Flechard, C.; Hamon, Y.; Busnot, S. (2019): 10 ans de mesures des concentrations atmosphériques NH₃ (gaz), NH₄⁺ (aérosol) et espèces chimiques associées sur sites agricoles et forestiers en France. colloque agriculture et qualité de l'air. INRA Paris, 2019.

Fenn, M. E.; Bytnerowicz, A.; Schilling, S. L.; Vallano, D. M.; Zavaleta, E. S.; Weiss, S. B. et al. (2018): On-road emissions of ammonia: An underappreciated source of atmospheric nitrogen deposition. In *Science of the Total Environment* 625, pp. 909–919. DOI: 10.1016/j.scitotenv.2017.12.313.

- Fowler, D.; Pilegaard, K.; Sutton, M. A.; Ambus, P.; Raivonen, M.; Duyzer, J. et al. (2009): Atmospheric composition change: Ecosystems–Atmosphere interactions. In *Atmospheric Environment* 43 (33), pp. 5193–5267. DOI: 10.1016/j.atmosenv.2009.07.068.
- Franzaring, J.; Holz, I.; Fangmeier, A. (2012): Effects of increased autumn temperatures and sub-acute levels of ammonia on post-winter development of four cultivars of winter oilseed rape (*Brassica napus* L.). In *Journal of Applied Botany and Food Quality* 85 (2), pp. 134–143.
- Frati, L.; Santoni, S.; Nicolardi, V.; Gaggi, C.; Brunialti, G.; Guttova, A. et al. (2007): Lichen biomonitoring of ammonia emission and nitrogen deposition around a pig stockfarm. In *Environmental pollution (Barking, Essex: 1987)* 146 (2), pp. 311–316. DOI: 10.1016/j.envpol.2006.03.029.
- Freiberger, L.; Schinkel, F.; Vogt, S.; Windisch, U. (2021): Kalibrierung von Bioindikationsverfahren zum Nachweis von Immissionen atmosphärischer reaktiver Stickstoffverbindungen/Calibration of bioindication methods for the detection of immissions of atmospheric reactive nitrogen compounds. In *GrdL* 81 (05-06), pp. 175–183. DOI: 10.37544/0949-8036-2021-05-06-21.
- García-Gómez, H.; Aguilauame, L.; Izquieta-Rojano, S.; Valiño, F.; Àvila, A.; Elustondo, D. et al. (2016): Atmospheric pollutants in peri-urban forests of *Quercus ilex*: evidence of pollution abatement and threats for vegetation. In *Environmental science and pollution research international* 23 (7), pp. 6400–6413. DOI: 10.1007/s11356-015-5862-z.
- Geßler, A.; Rienks, M.; Rennenberg, H. (2002): Stomatal uptake and cuticular adsorption contribute to dry deposition of NH₃ and NO₂ to needles of adult spruce (*Picea abies*) trees. In *New Phytologist* 156 (2), pp. 179–194. DOI: 10.1046/j.1469-8137.2002.00509.x.
- Husted, S.; Schjoerring, J. K. (1996): Ammonia Flux between Oilseed Rape Plants and the Atmosphere in Response to Changes in Leaf Temperature, Light Intensity and Air Humidity. Interactions with Leaf Conductance and Apoplastic NH₄⁺ and H⁺ Concentrations. In *Plant Physiology* 112, pp. 67–74. DOI: 10.1104/pp.112.1.67.
- Hutchinson, G. L.; Millington, R. J.; Peters, D. B. (1972) Atmospheric ammonia: Absorption by plant leaves. In *Science* 175:771–772.
- Ilugo Chibuzo, F. (2012): Biomonitoring of ammonia deposition by means of higher plants. Dissertation. University of Hohenheim, Stuttgart.
- Izquieta-Rojano, S.; López-Aizpún, M.; Irigoyen, J. J.; Santamaría, J. M.; Santamaría, C.; Lasheras, E. et al. (2018): Eco-physiological response of *Hypnum cupressiforme* Hedw. to increased atmospheric ammonia concentrations in a forest agrosystem. In *Science of the Total Environment* 619-620, pp. 883–895. DOI: 10.1016/j.scitotenv.2017.11.139.
- Jäger, H.-J.; Fangmeier, A.; Jong, M. de; Hanstein, S.; Pahlich, E. (1998): Untersuchung der Zusammenhänge zwischen Nährstoffeintrag und Vegetationsveränderung beim Halbtrockenrasen. Ableitung von ökologischen Schutzziele (FE-Vorhaben Nr. 108 03 050/01). Justus-Liebig-Universität Gießen.

Jones, L.; Nizam, M. S.; Reynolds, B.; Bareham, S.; Oxley, E.R.B. (2013): Upwind impacts of ammonia from an intensive poultry unit. In *Environmental Pollution* 180, pp. 221–228. DOI: 10.1016/j.envpol.2013.05.012.

Jones, M. R.; Leith, I. D.; Fowler, D.; Raven, J. A.; Sutton, M. A.; Nemitz, E. et al. (2007a): Concentration-dependent NH₃ deposition processes for mixed moorland semi-natural vegetation. In *Atmospheric Environment* 41 (10), pp. 2049–2060. DOI: 10.1016/j.atmosenv.2006.11.003.

Jones, M. R.; Leith, I. D.; Raven, J. A.; Fowler, D.; Sutton, M. A.; Nemitz, E. et al. (2007b): Concentration-dependent NH₃ deposition processes for moorland plant species with and without stomata. In *Atmospheric Environment* 41 (39), pp. 8980–8994. DOI: 10.1016/j.atmosenv.2007.08.015.

Jones, M. R.; Raven, J. A.; Leith, I. D.; Cape, J. N.; Smith, R. I.; Fowler, D. (2008): Short-term flux chamber experiment to quantify the deposition of gaseous ¹⁵N-NH₃ to *Calluna vulgaris*. In *Agricultural and Forest Meteorology* 148 (6-7), pp. 893–901. DOI: 10.1016/j.agrformet.2007.12.003.

Kelleghan, D. B.; Hayes, E. T.; Everard, M.; Keating, P.; Lesniak-Podsiadlo, A.; Curran, T. P. (2021): Atmospheric ammonia and nitrogen deposition on Irish Natura 2000 sites: Implications for Irish agriculture. In *Atmospheric Environment* 261 (1566–7197), p. 118611. DOI: 10.1016/j.atmosenv.2021.118611.

Krupa, S. V. (2003): Effects of atmospheric ammonia (NH₃) on terrestrial vegetation: a review. In *Environmental Pollution* 124 (2), pp. 179–221. DOI: 10.1016/S0269-7491(02)00434-7.

Leith, I. D.; van Dijk, N.; Pitcairn, C.E.R.; Sheppard, L. J.; Sutton, M. A. (2009): Standardised grasses as biomonitors of ammonia pollution around agricultural point sources. In M. A. Sutton (Ed.): *Atmospheric Ammonia: Detecting Emission Changes and Environmental Impacts*: Springer Netherlands, pp. 269–279. Available online at https://www.scopus.com/inward/record.uri?eid=2-s2.0-84885755620&doi=10.1007%2f978-1-4020-9121-6_16&partnerID=40&md5=0c6efe8363cc258b5645310d0bd44539.

Levy, P.; Dijk, N.; Gray, A.; Sutton, M.; Jones, M.; Leeson, S. et al. (2019): Response of a peat bog vegetation community to long-term experimental addition of nitrogen. In *Journal of Ecology* 107 (3), pp. 1167–1186. DOI: 10.1111/1365-2745.13107.

LfU (2019): Ammoniak-Immissionsmessungen in Bayern 2006 – 2017. Fortführung 2015-2017. Bayerisches Landesamt für Umwelt. Augsburg.

Lockyer, D. R.; Whitehead, D. C. (1986): The Uptake of Gaseous Ammonia by the Leaves of Italian Ryegrass. In *Journal of Experimental Botany* 37 (180), pp. 919–927. DOI: 10.1093/jxb/37.7.919.

Lohrengel, B.; Hainsch, A.; Klasmeier, E.; Dämmgen, U.; Mohr, K.; Wallasch, M. (2012): PASSAMMONI - Passivsammler-Messungen zur Erfassung der Ammoniak-Belastung in Niedersachsen. Zentrale Unterstützungsstelle Luftreinhaltung, Lärm und Gefahrstoffe - ZUS LLG. Hildesheim.

Lolkema, D. E.; Noordijk, H.; Stolk, A. P.; Hoogerbrugge, R.; van Zanten, M. C.; van Pul, W. A. J. (2015): The Measuring Ammonia in Nature (MAN) network in the Netherlands. In *Biogeosciences* 12 (16), pp. 5133–5142. DOI: 10.5194/bg-12-5133-2015.

Loubet, B.; Milford, C.; Sutton, M. A.; Cellier, P. (2001): Investigation of the interaction between sources and sinks of atmospheric ammonia in an upland landscape using a simplified dispersion-exchange model. In *Journal of Geophysical Research* 106, 24,183-24,195. DOI: 10.1029/2001JD900238.

Martin, N. A.; Ferracci, V.; Cassidy, N.; Hook, J.; Battersby, R. M.; Di Meane, E. A. et al. (2019): Validation of ammonia diffusive and pumped samplers in a controlled atmosphere test facility using traceable Primary Standard Gas Mixtures. In *Atmospheric Environment* 15, pp. 453–462. DOI: 10.1016/j.atmosenv.2018.11.038.

Mayer, W.; Pfefferkorn-Dellali, V.; Türk, R.; Dullinger, S.; Mirtl, M.; Dirnböck, T. (2013): Significant decrease in epiphytic lichen diversity in a remote area in the European Alps, Austria. In *Basic and Applied Ecology* 14 (5), pp. 396–403. DOI: 10.1016/j.baae.2013.05.006.

Morillas, L.; Roales, J.; Cruz, C.; Munzi, S. (2021): Resilience of Epiphytic Lichens to Combined Effects of Increasing Nitrogen and Solar Radiation. In *Journal of fungi* (Basel, Switzerland) 7 (5). DOI: 10.3390/jof7050333.

Munzi, S.; Cruz, C.; Branquinho, C.; Cai, G.; Faleri, C.; Parrotta, L. et al. (2020): More tolerant than expected: Taking into account the ability of *Cladonia portentosa* to cope with increased nitrogen availability in environmental policy. In *Ecological Indicators* 119 (5), p. 106817. DOI: 10.1016/j.ecolind.2020.106817.

Munzi, S.; Cruz, C.; Branquinho, C.; Pinho, P.; Leith, I. D.; Sheppard, L. J. (2014): Can ammonia tolerance amongst lichen functional groups be explained by physiological responses? In *Environmental Pollution* 187, pp. 206–209. DOI: 10.1016/j.envpol.2014.01.009.

Nemitz, E.; Sutton, M. A.; Wyers, G. P.; Jongejan, P. A. C. (2004): Gas-particle interactions above a Dutch heathland: I. Surface exchange fluxes of NH₃, SO₂, HNO₃ and HCl. In *Atmospheric Chemistry and Physics* 4, pp. 989–1005. DOI: 10.5194/acp-4-989-2004.

Niederhauser, B. (2017): Metrology for Ammonia in Ambient Air. Final publishable JRP Report. ENV55 MetNH₃, EURAMET. unpublished.

Noordijk, H.; Braam, M.; Rutledge-Jonker, S.; Hoogerbrugge, R.; Stolk, A. P.; van Pul, W. A. J. (2020): Performance of the MAN ammonia monitoring network in the Netherlands. In *Atmospheric Environment* 228. DOI: 10.1016/j.atmosenv.2020.117400.

Pan, Y.; Tian, S.; Wu, D.; Xu, W.; Zhu, X.; Liu, C. et al. (2020): Ammonia should be considered in field experiments mimicking nitrogen deposition. In *Atmospheric and Oceanic Science Letters* 13 (3), pp. 248–251. DOI: 10.1080/16742834.2020.1733919.

Paoli, L.; Benesperi, R.; Proietti Pannunzi, D.; Corsini, A.; Loppi, S. (2014): Biological effects of ammonia released from a composting plant assessed with lichens. In *Environmental Science and Pollution Research* 21 (9), pp. 5861–5872. DOI: 10.1007/s11356-014-2526-3.

Paoli, L.; Pirintsos, S.Arg.; Kotzabasis, K.; Pisani, T.; Navakoudis, E.; Loppi, S. (2010): Effects of ammonia from livestock farming on lichen photosynthesis. In *Environmental Pollution* 158 (6), pp. 2258–2265. DOI: 10.1016/j.envpol.2010.02.008.

Pinho, P.; Dias, T.; Cruz, C.; Sim Tang, Y.; Sutton, M. A.; Martins-Loução, M.-A. et al. (2011): Using lichen functional diversity to assess the effects of atmospheric ammonia in Mediterranean woodlands. In *Journal of Applied Ecology* 48 (5), pp. 1107–1116. DOI: 10.1111/j.1365-2664.2011.02033.x.

Pinho, P.; Llop, E.; Ribeiro, M. C.; Cruz, C.; Soares, A.; Pereira, M. J.; Branquinho, C. (2014): Tools for determining critical levels of atmospheric ammonia under the influence of multiple disturbances. In *Environmental Pollution* 188, pp. 88–93. DOI: 10.1016/j.envpol.2014.01.024.

Pinho, P.; Theobald, M. R.; Dias, T.; Tang, Y. S.; Cruz, C.; Martins-Loução, M. A. et al. (2012): Critical loads of nitrogen deposition and critical levels of atmospheric ammonia for semi-natural Mediterranean evergreen woodlands. In *Biogeosciences* 9 (3), pp. 1205–1215. DOI: 10.5194/bg-9-1205-2012.

Pitcairn, C.E.R.; Leith, I. D.; Sheppard, L.; Sutton, M. A. (1998): The Relationship between Nitrogen Deposition, Species Composition and Foliar Nitrogen Concentrations in Woodland Flora in the Vicinity of Livestock Farms. In *Environmental Pollution* 102, pp. 41–48. DOI: 10.1016/S0269-7491(98)80013-4.

Pogány, A.; Balslev-Harder, D.; Braban, C. F.; Cassidy, N.; Ebert, V.; Ferracci, V. et al. (2016): A metrological approach to improve accuracy and reliability of ammonia measurements in ambient air. In *Measurement Science and Technology* 27 (11). DOI: 10.1088/0957-0233/27/11/115012.

Puchalski, M. A.; Sather, M. E.; Walker, J. T.; Lehmann, C. M. B.; Gay, D. A.; Mathew, J.; Robarge, W. P. (2011): Passive ammonia monitoring in the United States: comparing three different sampling devices. In *Journal of environmental monitorin*: JEM 13 (11), pp. 3156–3167. DOI: 10.1039/c1em10553a.

Reinds, G. J.; Bak, J.; Rouil, L.; Scheuschner, T.; Schaap, M.; Hendriks, C. et al. (2019): Ammonia Regulations in Northern Europe. Summary of policies and practices in France, Germany, the United Kingdom, the Netherlands and Denmark Scientific Report No. 321. Available online at <http://dce2.au.dk/pub/SR321.pdf>.

Seitler, E.; Meier, M. (2022): Ammoniakimmissionsmessungen in der Schweiz 2000-2021, Messbericht, Forschungsstelle für Umweltbeobachtung FUB. Available online at https://www.bafu.admin.ch/dam/bafu/de/dokumente/luft/externe-studien-berichte/ammoniak-immissionsmessungen-in-der-schweiz-2000-2021.pdf.download.pdf/Ammoniak-Immissionsmessungen_CH_2000-2021.pdf

Sheppard, L. J.; Leith, I. D.; Mizunuma, T.; Neil Cape, J.; Crossley, A.; Leeson, S. et al. (2011): Dry deposition of ammonia gas drives species change faster than wet deposition of ammonium ions: Evidence from a long-term field manipulation. In *Global Change Biology* 17 (12), pp. 3589–3607. DOI: 10.1111/j.1365-2486.2011.02478.x.

Suarez-Bertoa, R.; Astorga, C. (2016): Isocyanic acid and ammonia in vehicle emissions. In *Transportation Research Part D: Transport and Environment* 49, pp. 259–270. DOI: 10.1016/j.trd.2016.08.039.

Sutton, M. A.; Pitcairn, C.E.R.; Fowler, D. (1993): The Exchange of Ammonia Between the Atmosphere and Plant Communities. In *Advances in Ecological Research* 24, pp. 301–393. DOI: 10.1016/S0065-2504(08)60045-8.

Sutton, M. A.; van Dijk, N.; Levy, P. E.; Jones, M. R.; Leith, I. D.; Sheppard, L. J. et al. (2020): Alkaline air: changing perspectives on nitrogen and air pollution in an ammonia-rich world. In *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences* 378 (2183), p. 20190315. DOI: 10.1098/rsta.2019.0315.

Tang, Y. S.; Braban, C. F.; Dragosits, U.; Dore, A. J.; Simmons, I.; van Dijk, N. et al. (2018): Drivers for spatial, temporal and long-term trends in atmospheric ammonia and ammonium in the UK. In *Atmospheric Chemistry and Physics* 18 (2), pp. 705–733. DOI: 10.5194/acp-18-705-2018.

Temple, P. J.; Harper, D. S.; Pearson, R. G.; Linzon, S. N. (1979): Toxic effects of ammonia on vegetation in Ontario. In *Environmental Pollution* 20 (4), pp. 297–302. DOI: 10.1016/0013-9327(79)90152-6.

Thöni, L.; Seitler, E.; Meier, M.; Kosonen, Z. (2018): Ammoniak-Immissionsmessungen in der Schweiz 2000-2018. Messbericht. Im Auftrag des Bundesamtes für Umwelt (BAFU), der OSTLUFT, der ZUDK, des Fürstentums Liechtenstein und der Kantone AG, AI, BE, BL/BS, FR, GL, GR, LU, NE, SG, SH, SO, TG, ZG und ZH. Forschungsstelle für Umweltbeobachtungen. Available online at <https://www.bafu.admin.ch>.

Twigg, M. M.; Berkhout, A. J. C.; Cowan, N.; Crunaire, S.; Dammers, E.; Ebert, V. et al. (2022): In-situ measurements of NH₃: instrument performance and applicability. In *Atmospheric Measurement Techniques* [preprint]. DOI: 10.5194/amt-2022-107.

Umweltbundesamt (2018): FORESTFLUX – Standörtliche Validierung der Hintergrunddeposition reaktiver Stickstoffverbindungen. Umweltbundesamt. Dessau-Roßlau.

Umweltbundesamt (2021): Ammoniak-Emissionen. Umweltbundesamt. Dessau-Roßlau. Available online at <https://www.umweltbundesamt.de/daten/luft/luftschadstoff-emissionen-in-deutschland/ammoniak-emissionen#entwicklung-seit-1990>, checked on 2/9/2022.

UNECE (2007): Review of the 1999 Gothenburg Protocol. Report on the Workshop on Atmospheric Ammonia: Detecting Emission Changes and Environmental Impacts. Available online at http://www.ammonia-ws.ceh.ac.uk/documents/ece_eb_air_wg_5_2007_3_e.pdf, checked on 9/2/2022.

van der Eerden, L. J. (1982): Toxicity of ammonia to plants. In *Agriculture and Environment* 7 (3-4), pp. 223–235. DOI: 10.1016/0304-1131(82)90015-7.

van der Eerden, L. J.; Dueck, Th. A.; Berdowski, J. J. M.; Greven, H.; van Dobben, H. F. (1991): Influence of NH₃ and (NH₄)₂SO₄ on heathland vegetation. In *Acta Botanica Neerlandica* 40 (4), pp. 281–296. DOI: 10.1111/j.1438-8677.1991.tb01559.x.

van Dijk, N.; Leeson, S.; Jones, M.; Sheppard, L.; Leith, I. (2017): What does adding Nitrogen to a bog do to the soil water? CEH Edinburgh, 2017. Available online at <https://www.caper.ceh.ac.uk/sites/default/files/Netty%20van%20Dijk.pdf>, checked on 2/9/2022.

van Hove; L. W. A.; Koops, A. J.; Adema, E. H.; Vredenberg, W. J.; Pieters, G. A. (1987): Analysis of the uptake of atmospheric ammonia by leaves of *Phaseolus vulgaris* L. In *Atmospheric Environment* 21 (8), pp. 1759–1763. DOI: 10.1016/0004-6981(87)90115-6.

VMM (2017): Tijdelijk meetnet ammoniak in Natura 2000-gebieden. eindrapport. Vlaamse Milieumaatschappij. Afdeling Lucht, Aalst.

Volten, H.; Bergwerff, J. B.; Haaima, M.; Lolkema, D. E.; Berkhout, A. J. C.; van der Hoff, G. R. et al. (2012a): Two instruments based on differential optical absorption spectroscopy (DOAS) to measure accurate ammonia concentrations in the atmosphere. In *Atmospheric Measurement Techniques* 5 (2), pp. 413–427. DOI: 10.5194/amt-5-413-2012.

Volten, H.; Haaima, M.; Swart, D.P.J.; van Zanten, M. C.; van Pul, W.A.J. (2012b): Ammonia exchange measured over a corn field in 2010. RIVM. Bilthoven.

Watmough, S. A.; McDonough, A. M.; Raney, S. M. (2014): Characterizing the influence of highways on springtime NO₂ and NH₃ concentrations in regional forest monitoring plots. In *Environmental pollution* (Barking, Essex: 1987) 190, pp. 150–158. DOI: 10.1016/j.envpol.2014.03.023.

Weldon, J.; Grandin, U. (2021): Weak recovery of epiphytic lichen communities in Sweden over 20 years of rapid air pollution decline. In *The Lichenologist* 53 (2), pp. 203–213. DOI: 10.1017/S0024282921000037.

Whitehead, D. C.; Lockyer, D. R. (1987): The Influence of the Concentration of Gaseous Ammonia on its Uptake by the Leaves of Italian Ryegrass, with and without an Adequate Supply of Nitrogen to the Roots. In *Journal of Experimental Botany* 38 (190), pp. 818–827. DOI: 10.1093/jxb/38.5.818.

Wichink Kruit, R. J.; Aben, J.; Vries, W. de; Sauter, F.; van der Swaluw, E.; van Zanten, M. C.; van Pul, W.A.J. (2017): Modelling trends in ammonia in the Netherlands over the period 1990–2014. In *Atmospheric Environment* 154, pp. 20–30. DOI: 10.1016/j.atmosenv.2017.01.031.

Wuytack, T.; Verheyen, K.; Wuyts, K.; Adriaenssens, S.; Staelens, J.; Samson, R. (2013): The use of leaf characteristics of common oak (*Quercus robur* L.) to monitor ambient ammonia concentrations. In *Water, Air, and Soil Pollution* 224 (1). DOI: 10.1007/s11270-012-1356-5.

Yao, X. H.; Zhang, L. (2013): Analysis of passive-sampler monitored atmospheric ammonia at 74 sites across southern Ontario, Canada. In *Biogeosciences* 10 (12), pp. 7913–7925. DOI: 10.5194/bg-10-7913-2013.

Yao, X. H.; Zhang, L. (2016): Trends in atmospheric ammonia at urban, rural, and remote sites across North America. In *Atmospheric Chemistry and Physics* 16 (17), pp. 11465–11475. DOI: 10.5194/acp-16-11465-2016.

Yao, X. H.; Zhang, L. (2019): Causes of Large Increases in Atmospheric Ammonia in the Last Decade across North America. In *ACS omega* 4 (26), pp. 22133–22142. DOI: 10.1021/acsomega.9b03284.

3 Discussion points at the workshop

Before and during the workshop, we aimed to collate recent information on whether critical levels of ammonia need to be revised and asked the presenters and participants to notify the audience about ongoing research activities. In order to prepare a general discussion, we raised the following questions prior to the workshop. Participants were asked to respond before the meeting, but we did not get any feedback until the workshop. However, we were still able to address some of the points at the final discussion (see chapter 4.3)

Discussion 1: Current set-up and future perspective on ammonia monitoring

- With regard to ammonia measuring, is the use of passive samplers suitable? At which frequency and at how many sites should samplers be analysed?
- Are there new/suitable/affordable measuring devices to continuously monitor NH₃ in the low concentration range?
- Which criteria have to be met to set up a national representative ammonia network?
- Which difficulties complicate ammonia monitoring networks? Costs?
- Is there essential knowledge for recommendations to policy makers that is gained (exclusively) with ammonia monitoring networks?
- Is ammonia monitoring beneficial for nature protection and air quality policies?
- Would you support an ammonia monitoring obligation and for what reasons?

Discussion 2: New findings with potential relevance to NH₃ CLe: Future directions


- Do you perform or are you aware of research on ammonia effects on higher or lower plants using controlled fumigation at ammonia levels < 10 µg m⁻³? What are the technologies involved?
- Do you perform or do you know of ammonia fumigation studies in the field? What are the technologies involved and which are the tested ecosystems?
- Do you perform or do you know of gradient studies using active or passive biomonitoring, i.e. the use of vegetation on-site or plants being actively exposed? What are the involved technologies and plant species?
- Would you support annual, monthly and/or short-term CLe?
- Would you support to include knowledge on Critical Loads into scientific justification of critical levels?
- Based on current research, is there (enough) evidence for changing/keeping the CLe (for lower and higher plants, respectively)?
- Which further steps after the Workshop should be taken in the review of NH₃ CLe?

4 Workshop presentations and discussion

The workshop on CLe for ammonia took place as a hybrid meeting on 28/29 March 2022 with the option of physical attendance at the German Environment Agency in Dessau, Germany. 15 participants were able to attend in presence and over 100 people were following the live stream via a WebEx online meeting platform from several European countries and the US. In total, there were 145 registered participants from 28 countries including representatives from several bodies to the convention such as the UNECE Secretariat, WGE, ICP Vegetation, ICP Forest, CDM, CIAM, MSC-West, TFRN and representatives from several EU Environment Agencies. 19 presentations from presenters from nine countries (Figure 2) were dealing with a current review, models and future trends of NH₃ across Europe in the first session, different ammonia monitoring networks in the second session and with vegetation effects (recent research on different scales) in the third session.

The workshop started with a welcome message from Markus Geupel (UBA) introducing the aim and ideas of the workshop. As the last ammonia workshop took place in 2006 in Edinburgh, the first presentation dealt with a recap of the issues and topics back then. Then, a presentation about recent and first research on ammonia effects on vegetation was given, summarizing the most important points of a background document (updated version in Chapter 2 or present report). The rest of the presentations on day 1 addressed modelling and monitoring approaches in different countries.

Figure 2: Numbers of presentations per country



Source: own illustration

In the following, we present the workshop agenda and in the sub-chapters thereafter, we collate the written summaries which were provided by the presenters. Finally, we document the most important discussion points and come up with the key findings.

Expert Workshop on Ammonia

28.-29. March 2022

Held hybrid: German Environment Agency, Dessau / webex

All times are given in CEST

More than ten years after the recommendation of updated CLRTAP critical levels for ammonia ([Mapping Manual](#)), new findings on the effects of ammonia on vegetation have been discussed on a workshop prepared by CCE and Germany. Furthermore, the workshop offered the opportunity to exchange information on national or regional programs which have been set up for the monitoring of ammonia in sensitive habitats.

Scientists dealing with research on effects of ammonia on vegetation and ecosystems or monitoring ammonia in the environment were encouraged to present their research.

The expert workshop was held on 28 to 29 March 2022 as a hybrid meeting. The option for a physical attendance was in Dessau, Germany, where the German Environment Agency (UBA) is situated. In parallel, online participation was offered. The workshop was part of the current work plan for the year 2022-2023 of CLRTAP ([1.1.1.22](#)) and was organized in the framework of an R&D Project in Germany financed by the German Environment Agency (UBA).

FINAL AGENDA (Number of presentations on day 1 is the same as in the sub-chapters)

All presentations can be accessed under <https://clous.uba.de/index.php/s/Hk15ZLnlpF1ZWqb> with the following access code: Ammoniak@uba

Day 1 (Monday 28 March 2022)

12.30-13.50 Session 1: Introduction (Background document, Current Review, models and future trends of NH₃ Concentrations across Europe)

Introduction to the workshop and the review process (Geupel, UBA, Germany)

- 1. 15 years after: Rationale and approach used in the 2006 Edinburgh review workshop (Sutton, CEH, United Kingdom)**
- 2. Literature review on the effects of ammonia (Franzaring and Kösler, University of Hohenheim, Germany)**
- 3. Modelling ammonia concentrations, trends and scenarios in Europe (Fagerli, MSC-West, Norway)**

13.50 - 15.10 Session 2: Ammonia monitoring networks (national, regional, in conservation areas)

- 4. NH₃ concentration measurements (Bleeker, RIVM, The Netherlands)**
- 5. Real-time monitoring of ammonia concentrations with Cavity Ring-Down Spectroscopy' (Hofmann, Picarro, The Netherlands)**
- 6. Ammonia monitoring at long-term and temporary sites in Flanders (Staelens, Flanders Environment Agency, Belgium)**
- 7. Monitoring atmospheric ammonia on Natura 2000 sites in the Republic of Ireland (Kelleghan, University College Dublin, Ireland)**

15.10 -15.40 Break

15.40 - 17.30 Session 2 continued

- 8. UK Ammonia Network: present and future (Aazem, JNCC, United Kingdom)**
- 9. Ammonia concentration measurements in Germany - an overview (Moravek & Geupel, UBA, Germany)**
- 10. Metrological aspects to support environmental ammonia monitoring (Pogany, PTB, Germany)**
- 11. Long-term NH₃ measurements at the TROPOS research station Melpitz using a MARGA (Poulain, TROPOS, Germany)**
- 12. Long-term measurements to support deposition and critical loads assessments in the US (Gay, University of Wisconsin Madison & Puchalski, EPA, USA)**

17.30-18.00 Discussion 1: Current set-up and future perspectives on ammonia monitoring

Day 2 (Tuesday 29 March 2022)

9.00 – 10.40 Session 3: Recent research on different scales (fumigation chamber, field exposure, gradient studies with bioindicators, transplants and vegetation on site)

- 13. Relationships between ambient NH₃ concentrations and epiphytic lichens in an urban environment (Manninen, University of Finland, Finland)**
- 14. Results from an ammonia fumigation and a field study using endangered nitrogen-sensitive plant species (Kösler and Franzaring University of Hohenheim, Germany)**
- 15. Towards understanding shorter-term critical levels with monthly resolution measurements (Dragosits, CEH, United Kingdom)**
- 16. Biomonitoring of ammonia with the epiphytic lichen *Hypogymnia physodes* (Mohr, LWK Lower Saxony, Germany)**
- 17. How can we assess the impact of project contributions of ammonia on Natura 2000 sites? (Uhl, FÖA, Germany)**

10.40 – 11.00 Coffee Break

11.00-11.40 Session 4: Interrelations of critical levels and critical loads

- 18. Development of a trophic assessment and management system for habitat types (Directive 92/43/EWG) considering interrelations between critical levels and critical loads (Prüeß, LUBW Baden-Württemberg, Germany)**
- 19. Ammonia concentrations in Switzerland compared to critical levels and loads (Meier, BAFU, Switzerland)**

11.40 - 13.00 Discussion 2: New findings with potential relevance to NH₃ critical levels: Future directions / next steps

4.1 Workshop Day 1, 28 March 2022

4.1.1 15 years on: Rationale and reflection on the 2006 Edinburgh Ammonia Workshop in the light of emerging evidence.

Mark A. Sutton¹, Patricia A. Wolseley², Netty van Dijk¹, Ajinkya G. Deshpande¹, Ulrike Dragosits¹, Marsailidh Twigg¹, Y. Sim Tang¹, Christine Braban¹, Pete Levy¹, William J. Bealey¹, Gothamie Weerakoon², Jize Jiang³, David S. Stevenson³, Buddhika Weerakoon⁴, Sarath P. Nissanka⁴ and Matthew R. Jones¹

¹ UK Centre for Ecology & Hydrology, United Kingdom

² Natural History Museum, London

³ University of Edinburgh, School of GeoSciences

⁴ University of Peradeniya, Sri Lanka.

Background

A critical level for ammonia was first set in the 1980s at Bad Harzburg (Posthumus, 1988), which was subsequently reassessed in the 1992 Egham Workshop (Ashmore and Wilson, 1994). However, there were several arguments subsequently articulated for why further revision was needed. The original critical level included an annual value of 8 $\mu\text{g m}^{-3}$, but comparison with dry deposition rates showed that there was an inconsistency between critical loads and critical levels, with critical loads exceeded at much lower ammonia concentrations than critical levels. In addition, NH₃ monitoring data suggested that there was an increased likelihood of exceedance of the critical level for averaging periods of longer than one year (see further below, regarding Figure 3), while field data suggested responses to NH₃ concentrations in the field at much less than 8 $\mu\text{g m}^{-3}$ (Sutton et al., 2009a). It was with these issues in mind, that the UK government agreed to sponsor the 2006 Edinburgh Atmospheric Ammonia Workshop (CEH, 2006), which led to revision of the ammonia critical level (UNECE, 2007) and a comprehensive publication that also covered other aspects of atmospheric ammonia (Sutton et al., 2009b). The workshop itself took place at the Scottish Government offices, with 80 experts from 19 parties to the UNECE Convention on Long-Range Transboundary Air Pollution, including support from COST729, Defra, UK Centre for Ecology & Hydrology, Scottish Government and the NitroEurope Integrated Project (CEH, 2006).

The following sections summarize key objectives of the workshop and then reflect on the developing picture, in particular considering evidence that has emerged over the last 15 years since the Edinburgh review.

Objective 1: Review of the ammonia critical level

The first objective was to assess the extent to which the existing critical thresholds for ammonia reflect current scientific understanding (Sutton et al., 2009b, p 4):


- a) To examine the case for setting new ammonia critical threshold(s) based on current evidence of direct impacts of ammonia on different receptors;
- b) To discuss the extent of differential sensitivity to ammonia versus other forms of reactive N;
- c) To debate the case for establishing indicative air concentration limits for indirect effects of ammonia, which would be consistent with current critical loads for nitrogen.

It was concluded that there was substantial evidence of long-term impacts of ammonia, such as from the Whim Bog experiment, running at that time already over 6 years, and now running for around 20 years. The most recent analyses (Levy et al., 2019; Sutton et al., 2020), emphasize that ammonia impacts integrate over periods of much longer than one year, but are also faster in occurring than the effects of equivalent amounts of wet deposited nitrogen. This provides long-term evidence that gaseous ammonia is more damaging to some forms of vegetation than wet deposited nitrogen, highlighting the importance of recognizing these impacts. It still needs to be determined if there is sufficient data available to revise critical loads based on current understanding for the differing sensitivity of dry deposited gaseous ammonia versus wet deposited nitrogen. Similarly, it was not agreed in the Edinburgh Workshop to establish indicative air concentration limits by simple equivalence to critical loads. One of the concerns here is uncertainty in dry deposition rates, especially when considering bi-directional fluxes. By contrast, adoption of the revised critical levels for ammonia has given appropriate prominence to its adverse effects.

One of the outcomes from the Edinburgh Workshop that has received least attention is the demonstrated relationship between the previous critical level and long-term ammonia monitoring data (Figure 3). This showed that the 1992 critical levels are most precautionary over longer averaging periods, compared with shorter averaging periods (Sutton et al., 2009c). Hence, in Figure 3 it can be seen that the old annual critical level is more likely to be exceeded than the hourly and daily critical levels for the NH₃ monitoring datasets examined. Again, this demonstrated the importance of setting a long-term critical level (expressed as an annual mean, but designed for application over several decades).

Figure 3: Comparison of ammonia concentration data from long term monitoring with the Egham 1992 critical levels for ammonia, as well as an extrapolated potential critical level for averaging over 25 years.

The data from multiple sites show that the critical level is most precautionary over longer averaging periods.



Source: Sutton et al., 2009c

The previous UNECE Critical Levels from the Egham Workshop 1992 were as follows:

- 8 $\mu\text{g m}^{-3}$ annual mean,
- 23 $\mu\text{g m}^{-3}$ monthly mean,
- 270 $\mu\text{g m}^{-3}$ daily mean,
- 3300 $\mu\text{g m}^{-3}$ hourly mean.

The further evidence from multiple field studies including from the UK, Switzerland, Portugal and Italy (Cape et al., 2009a,b) showed that the annual mean value was not sufficiently precautionary, and that effects accumulate for periods longer than 1 year. Data from the Netherlands were also available, but it was concluded that ambient ammonia concentrations were too high to derive a critical level from these data.

The following values were agreed in Edinburgh (Sutton et al., 2009b; Cape et al., 2009a,b) and subsequently included in the UNECE Mapping Manual (current edition: UNECE, 2017):


‘Long-term’ annual average critical level for lichens, bryophytes and for ecosystems in which they are important were set at 1 $\mu\text{g NH}_3 \text{ m}^{-3}$.

‘Long-term annual average critical level for higher vegetation (inc. woodland ground flora): were set at 3 [2-4] $\mu\text{g NH}_3 \text{ m}^{-3}$.

These critical levels were set as the long-term values, which cannot be assumed to protect for >20-30 years (expressed as annual mean). Conversely, it was concluded that there was insufficient evidence to revise the short-term critical level values (monthly, daily, hourly).

Figure 4: Possible mechanisms by which atmospheric NH_3 pollution affects epiphytic lichens, including both positive (+) and negative (-) effects.

Solid lines indicate observed relationships or those directly implied by physico-chemistry.



Source: Sutton et al., 2020


Subsequent analysis since 2006 has given further understanding of the potential mechanisms for why ammonia should have a more adverse effect than ammonium. This includes the observation that ammonia has an ‘alkaline air’ effect, indicated by the observation that the pH of tree bark and

other vegetation surfaces increases in the vicinity of ammonia sources (Figure 4). The relative importance of nitrogen eutrophication vs the alkaline air impacts requires further investigation.

Further evidence that needs consideration suggests that NH_3 and NO_x can have combined effects. Analysis of UK-wide data for epiphytic lichens (Lewis, 2012) identified a relationship with Lichen Indicator Score (LIS) leading to a Nitrogen Air Quality Index (NAQI), which has been applied by the UK Field Studies Council in its lichen nitrogen air quality guide (Wolseley et al. 2014). The following relationship was identified based on the UK data, and applied in Figure 5.

$$\text{NAQI: Nitrogen Air Quality Index} = 2[\text{NH}_3]^{0.5} + [\text{NO}_2]^{0.5} \quad (\text{using } \mu\text{mol m}^{-3}).$$

Figure 5: Relationship between the Lichen Indicator Score (LIS) and the Nitrogen Air Quality Index (NAQI).



Source: Wolseley et al., 2013

Based on this NAQI relationship derived from the UK epiphytic lichen data, the following scenarios can be envisaged:

$1 \mu\text{g m}^{-3} \text{NH}_3$ and $0 \mu\text{g m}^{-3} \text{NO}_2$	$\text{NAQI} = 0.49$	(Scenario 1)
$0 \mu\text{g m}^{-3} \text{NH}_3$ and $30 \mu\text{g m}^{-3} \text{NO}_2$	$\text{NAQI} = 0.8$	(Scenario 2)
$0 \mu\text{g m}^{-3} \text{NH}_3$ and $11.5 \mu\text{g m}^{-3} \text{NO}_2$	$\text{NAQI} = 0.5$	(Scenario 3)
$1 \mu\text{g m}^{-3} \text{NH}_3$ and $30 \mu\text{g m}^{-3} \text{NO}_2$	$\text{NAQI} = 0.49 + 0.8 = 1.29$	(Scenario 4)
$0.5 \mu\text{g m}^{-3} \text{NH}_3$ and $10 \mu\text{g m}^{-3} \text{NO}_2$	$\text{NAQI} = 0.34 + 0.47 = 0.81$	(Scenario 5).

Considering the scenarios outlined above, Scenario 1 shows the value of NAQI at the ammonia critical level, which links to the change from clean to at risk in Figure 5. By comparison, Scenario 2 is the standard NO_2 critical level ($\text{NAQI}=0.8$), which, according to the lichen epiphyte data, is not as precautionary as the NH_3 critical level ($\text{NAQI}=0.5$). Scenario 3 shows that the same level of precaution as the NH_3 critical level ($\text{NAQI}=0.5$) would be NO_2 of around $11.5 \mu\text{g m}^{-3}$. Scenario 4 illustrates that if both NH_3 and NO_2 are at the critical level, the overall threat to epiphytic lichens is larger ($\text{NAQI}=1.29$) than if only one gas is considered. Finally, Scenario 5 indicates a case where both NH_3 and NO_2 individually are less than the critical level, but the combined risk to epiphytic lichens ($\text{NAQI}=0.81$) is larger than having NH_3 alone at the critical level ($\text{NAQI}=0.5$).

These simple scenarios have implications for future consideration of how the NH_3 and NO_2 critical levels fit together. According to the UK data from which this relationship was derived, the NH_3 and

NO₂ effects are additive. Hence this suggests that a smaller critical level for NO₂ would be expected where NH₃ concentrations are high, and *vice versa*. This requires further investigation to inform future development of the critical levels approach for NH₃ and NO₂.

Objective 2: Atmospheric ammonia measurements

The second objective of the Edinburgh Workshop was to assess the extent to which independent atmospheric measurements can verify where regional changes in NH₃ emissions have and have not occurred (Sutton et al., 2009b, p 3):

- a) To quantify the extent to which regional changes in ammonia emissions have been reflected in measurements of ammonia and ammonium in the atmosphere;
- b) To distinguish cases where the estimated changes are due to altered sectoral activity or the implementation of abatement policies to see if atmospheric measurements verify the ammonia abatement policies;
- c) To make recommendations for future air monitoring and systems for assessing the national implementation of ammonia abatement policies and consider the implications of any non-linearities.

The overall state of the evidence was summarized by Bleeker et al. (2009), showing that an ‘ammonia gap’ in the Netherlands - between measured and modelled NH₃ concentrations - was partly explained. Trends over time were consistent, while differences in absolute values could be attributed to typical model-measurement differences. The differences between modelled versus measured trends could also be understood by considering the case of Hungary where ammonia concentrations had not decreased despite reductions in fertilizer use and animal numbers. Comparison with aerosol and wet deposition data and a modelling scenario showed that this could be understood because SO₂ emissions had declined over the period, leading to a longer atmospheric residence time for gaseous NH₃ (Horvath et al., 2009). The implication is that ammonia monitoring needs to consider all the phases, including gaseous, aerosol and wet deposition. Subsequent reports of 20-year time series of ammonia monitoring compared with models (e.g. van Zanten et al., 2017; Wichink Kruit, 2017; Tang et al., 2018) have further improved our understanding of the relationships between ambient ammonia concentrations and changing air chemistry. In parallel, Europe-wide measurements have shown the importance of ammonia and ammonium in the overall inorganic pollution load (Tang et al., 2020).

Objective 3: Ammonia Hotspots


The third objective of the Edinburgh Workshop was to review approaches for downscaling transboundary assessments to deal with ammonia ‘hotspots’ (locations in landscapes with very high ammonia levels) in relation to operational modelling and monitoring (Sutton et al., 2009b, p 5):

- a) To review current emission and atmospheric dispersion modelling methods for downscaling NH₃ dispersion and deposition in hotspots;
- b) To examine the status of methods for effect assessment and air monitoring in NH₃ hotspots;
- c) To recommend broad principles for assessment of ammonia hotspots, including interactions between transboundary ammonia emission reduction targets and other policy measures.

A major review for the workshop of the challenges of assessing nitrogen in hotspots was provided by Loubet et al. (2009). Assessing ammonia in hotspots remains an issue of high uncertainty,

especially emissions and dry deposition. The workshop concluded that there is a need for better knowledge on ammonia compensation points, surface resistances, and dependence on climatic variables and deposition history. It was noted that there is potential for local “landscape models” to develop local “tailored” abatement measures, as well as to quantify sub-grid variability (see Figure 6). Detailed landscape case studies have been provided by Dragosits et al. (2006), Cellier et al. (2011) and Vogt et al. (2013). Subsequent work has continued to highlight the fine scale spatial variability in ammonia concentrations, and the consequent risk for protected areas such as the Natura 2000 network (Hallsworth et al., 2010), as well as landscape-based mitigation opportunities (such as tree-belts and atmospheric buffer zones). In this case, enhanced rates of NH_3 deposition to woodland vegetation are used in landscape planning measures to increase deposition to targeted sites, e.g. non-conservation woodland, in order to help protect priority conservation sites (Bealey et al., 2016; Dalgaard et al., 2022).

Figure 6: Illustration of the challenges of dealing with spatial scale in nitrogen deposition and critical loads exceedances. The landscape case study is taken from.



Source: Dragosits et al. (2007)

Objective 4: Mesoscale modelling of ammonia

This final objective of the Edinburgh Workshop focused on reviewing mesoscale atmospheric transport and chemistry models in relation to their formulation and results for ammonia (Sutton et al., 2009c, p7):


- a) To review emission parameterizations used in the models, establishing comparability, spatial and temporal resolution and uncertainties;
- b) To review dispersion, air chemistry and deposition formulations identifying key differences and uncertainties;
- c) To assess the overall performance of the models against measurements, giving model recommendations and implications of non-linearities for integrated assessment models.

A detailed review was provided for the workshop by van Pul et al. (2009). Six models from national to regional (European) scale were reviewed. This showed that estimates of wet deposition and aerosol NH_4^+ concentrations are generally adequate. It was concluded that all European scale models currently underestimate ground level NH_3 concentrations, but better agreement is found in

national models. Key uncertainties were identified linked to emissions, dry deposition, spatial resolution and description of vertical diffusion. It was noted that none of the models routinely included compensation points, and a more comprehensive database was needed. This has since partly been provided by the review of Massad et al. (2010), though global scale extension is needed as evidence was predominantly from European and North American contexts. In addition to the emergence of satellite data as a key tool for monitoring ammonia trends (e.g. van Damme et al., 2018, 2021), perhaps the most significant message to be emphasized since this time is the strong climatic dependence of ammonia emissions. To date ammonia emissions inventories almost universally assume that emissions are not dependent on climate, so that emission factors are not affected by climate change. However, as illustrated by Sutton et al. (2013), Riddick et al. (2018) and Jiang et al. (2021) there is a substantial climate dependence where a warmer world is projected to emit more ammonia.

The case is clearly illustrated for poultry which excrete uric acid, where both temperature and water availability affect hydrolysis to form ammonia, and these same factors, together with windspeed also affect volatilization rates. The combined impact of these factors is shown in Figure 7 (Jiang et al. 2006). According to this mechanism NH₃ emissions could easily increase by around 30-40% over the 21st century solely due to climatic warming (Sutton et al., 2013; Riddick et al., 2018). This is of major importance in emphasizing the necessity of emission controls to avoid increasing ammonia levels and worsening the impacts on ecosystems (and human health) in future.

Figure 7: Simulated a) annual global NH₃ emissions (Gg N yr⁻¹) from poultry farming in 2010. b) Percentage of excreted nitrogen that volatilizes (PV, %) as NH₃ from chicken housing in 2010. The resolution is 0.5° 0.5°.



Source: Jiang et al. (2006)

Outlook

The Edinburgh Workshop provided a strong foundation for revising the ammonia critical level, which in the case of lichens, bryophytes and ecosystems where these are important to ecosystem integrity, reduced the critical level by a factor of eight (from 8 to 1 $\mu\text{g m}^{-3}$). Other evidence presented at the workshop has helped inform better understanding of ammonia monitoring, hotspot analysis and modelling at regional scales.

Since the Edinburgh Workshop, ongoing long-term experimentation has continued to demonstrate the sensitivity of vegetation to ammonia. Regional lichen surveying across the UK has also indicated the potential for an additive effect of NH_3 and NO_2 on lichens, which requires further investigation in the context of reviewing the NO_2 critical level.

There is an obvious need for further experimentation on the impacts of gaseous ammonia. In this context, the authors are grateful to UKRI for funding the GCRF South Asian Nitrogen Hub, which has recently established two new ammonia enrichment facilities in woodland ecosystems, one in Scotland and the other in Sri Lanka. These sites are expected to deliver improved understanding on ammonia critical levels in the future, including their effect on epiphytic lichens and bryophytes. Preliminary data from the Sri Lanka field site hosted by the Dilmah Conservation Center for Climate Change Research & Adaptation (DCC) (Queensbury Tea Estate), show that the site is highly suitable for the long-term monitoring experiment. Results showed the ammonia concentrations range from 0.4 $\mu\text{g m}^{-3}$ NH_3 in the natural forest (Rilagala forest) to 1.2 $\mu\text{g m}^{-3}$ at the edge of the forest (meteorological station at DCC) and 2.1 $\mu\text{g m}^{-3}$ in surrounding tea plantations, as compared with 2.4 $\mu\text{g m}^{-3}$ at a potato farm (Seetha Eliya). The datasets that will emerge from such new experiments are especially relevant in regions such as South Asia (Ellis et al., 2021), where limited evidence has until now been available compared with other regions.

Acknowledgements

The Whim Bog facility and research is supported by the NERC UK Status, Change and Projections of the Environment (UKSCAPE) National Capability Programme (NE/R016429/1). The work is also supported by UK Research and Innovation (UKRI) as a Global Challenges Research Fund grant, the 'GCRF South Asian Nitrogen Hub' (NE/S009019/1).

References

Ashmore M.R. and Wilson R.B. (1994) *Critical levels of air pollutants for Europe*. Report of the UNECE Expert Workshop at Egham, UK, 23-26 March 1992). Air Quality Division, Department of the Environment, London.

Bealey W.J., Dore A.J., Dragosits U., Reis S., Reay D.S. and Sutton M.A. (2016) The potential for tree planting strategies to reduce local and regional ecosystem impacts of agricultural ammonia emissions *Journal of Environmental Management* **165**, 106-116.

Bleeker A., Sutton M.A., Acherman B., Alebic-Juretic A., Aneja V.P., Ellermann T., Erisman J.W., Fowler D., Fagerli H., Gauger T., Harlen K.S., Hole L.R., Horváth L., Mitisinkova M., Smith R.I., Tang Y.S., van Pul A. (2009) Linking ammonia emission trends to measured concentrations and deposition of reduced nitrogen at different scales. Chapter 11, in: *Atmospheric Ammonia: Detecting emission changes and environmental impacts* (eds. M.A. Sutton, S. Reis and S.M.H. Baker), pp 123-180, Springer.

Cape J.N., van der Eerden L.J., Sheppard L.J., Leith I. D. and Sutton M.A. (2009a) Reassessment of critical levels for atmospheric ammonia. Chapter 2 in: *Atmospheric Ammonia, Detecting emission changes and environmental impacts*. pp 93-99. (eds. M.A. Sutton, S. Reis and S.M.H. Baker), Springer.

Cape J.N., van der Eerden L.J., Sheppard L.J., Leith I.D., Sutton M.A. (2009b) Evidence for changing the critical level for ammonia. *Environmental Pollution* **157**, 1033-1037. DOI: 10.1016/j.envpol.2008.09.049.

CEH (2006) Expert Workshop under the Convention on Long-range Transboundary Air Pollution Atmospheric Ammonia. Detecting emission changes and environmental impacts. <http://www.ammonia-ws.ceh.ac.uk/index.html> (last accessed 29 June 2022).

Cellier P., Bleeker A., Breuer L., Dalgaard T., Dragosits U., Drouet J.L., Durand P., Duret S., Hutchings N., Kros J., Loubet B., Oenema O., Olesen J.E., Mérot P., Theobald M.R., Viaud V., de Vries W. and Sutton M.A. Dispersion and fate of nitrogen in rural landscapes. Chapter 11, in: *The European Nitrogen Assessment* (Eds. Sutton M.A., Howard C.M., Erisman J.W., Billen G., Bleeker A., Grennfelt P., van Grinsven H. and Grizzetti B.) pp 229-248. Cambridge University Press.

Dalgaard T., Butterbach-Bahl K., Durand P., Hansen B., Lukin S., Moklyachuk L., Pelster D., Valinia S., Skiba U.M. and Sutton M.A. (2022) Land-use and landscape management. Chapter 6 in: *Nitrogen opportunities for agriculture, food & environment*. UNECE Guidance Document on Integrated Sustainable Nitrogen Management. UK Centre for Ecology & Hydrology, Edinburgh, UK (in press).

Dragosits U., Theobald M.R., Place C.J., ApSimon H.M. and Sutton M.A. (2006) The potential for spatial planning at the landscape level to mitigate the effects of atmospheric ammonia deposition. *J. Environ. Sci. & Policy* **9**, 626-638.

Ellis C.J., Steadman C.E., Vieno M., Sudipto C., Jones M.R., Negi S., Pandey B.P., Rai H., Tshering D., Weerakoon G., Wolseley P., Reay D., Sharma S. and Sutton M.A. (2022) Estimating nitrogen risk to Himalayan forests using thresholds for lichen bioindicators. *Biological Conservation* **265**, 109401. DOI: 10.1016/j.biocon.2021.109401.

Hallsworth S., Dore A.J., Bealey W.J., Dragosits U., Vieno M., Hellsten S., Tang Y.S. and M.A. Sutton M.A. (2010) The role of indicator choice in quantifying the threat of atmospheric ammonia to the 'Natura 2000' network. *Environmental Science and Policy* **13**, 671-687.

Horváth L., Fagerli H. and Sutton M.A. (2009) Long-Term Record (1981–2005) of ammonia and ammonium concentrations at K-Pusztá Hungary and the effect of sulphur dioxide emission change on measured and modelled concentrations. Chapter 12, in: *Atmospheric Ammonia: Detecting emission changes and environmental impacts* (eds. M.A. Sutton, S. Reis and S.M.H. Baker), pp 181-186, Springer.

Jiang J., Stevenson D.S., Uwizeye A., Tempio G. and Sutton M.A. (2021) A climate-dependent global model of ammonia emissions from chicken farming. *Biogeosciences* **18**, 135-158. <https://doi.org/10.5194/bg-18-135-2021>

Levy P., van Dijk N., Gray A., Sutton M., Jones M., Leeson S., Dise N., Leith I.D. and Sheppard L.J. (2019) Response of a peat bog vegetation community to long-term experimental addition of nitrogen. *J. Ecology* **107** (3), 1167-1186. DOI: 10.1111/1365-2745.13107.

Lewis J.E.J. (2012) Bio-monitoring for atmospheric nitrogen pollution using epiphytic lichens and bryophytes. PhD thesis, University of Nottingham. 297 pp. Available at: http://eprints.nottingham.ac.uk/13573/1/J_Lewis_PhD_thesis.pdf (last checked 29 June 2022).

Loubet B., Asman W.A.H., Theobald M.R., Hertel O., Tang Y.S., Robin P., Hassouna M., Daemmgen U., Genermont S., Cellier P. and Sutton M.A. (2009) Ammonia deposition near hot spots: processes, models and monitoring methods. Chapter 15, in: *Atmospheric Ammonia: Detecting emission changes and environmental impacts* (eds. M.A. Sutton, S. Reis and S.M.H. Baker), pp 205-267, Springer.

Massad R.-S., Nemitz E. and Sutton M.A. (2010) Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere *Atmos. Chem. Phys.* **10**, 10359-10386. www.atmos-chem-phys.net/10/10359/2010/

Posthumus A.C. (1988) Critical levels for effects of ammonia and ammonium. In: *Proceedings of the Bad Harzburg Workshop*. 117-127, UBA, Berlin.

Riddick, S.N., Dragosits, U., Blackall, T.D., Tomlinson, S.J., Daunt, F., Wanless, S., Hallsworth, S., Braban, C.F., Tang, Y.S. and Sutton, M.A. (2018) Global assessment of the effect of climate change on ammonia emissions from seabirds. *Atmos. Environ.* **184**, 212-223. <https://doi.org/10.1016/j.atmosenv.2018.04.038>

Sutton M.A., Reis R., Riddick S.N., Dragosits U., Nemitz E., Theobald M.R., Tang Y.S., Braban C.F., Vieno M., Dore A.J., Mitchell R.F., Wanless S., Daunt F., Fowler D., Blackall T.D., Milford. C., Flechard C.F., Loubet B., Massad R., Cellier P., Coheur P.F., Clarisse L., van Damme M., Ngadi, Y., Clerbaux C., Skjøth C.A., Geels C., Hertel O., Wichink Kruit R.J., Pinder, R.W., Bash J.O., Walker J.D., Simpson D., Horvath, L., Misselbrook, T.H., Bleeker A., Dentener F. & Wim de Vries V. (2013) Toward a climate-dependent paradigm of ammonia emission & deposition. *Phil. Trans. Roy. Soc. (Ser. B)*. 368: 20130166. <http://dx.doi.org/10.1098/rstb.2013.0166>

Sutton M.A., Reis S. and Baker S.M.H. (2009b) *Atmospheric Ammonia, Detecting emission changes and environmental impacts*. (eds.), Springer. 464 pp.

Sutton M.A., Sheppard L.J. and Fowler D. (2009a) Potential for the further development and application of critical levels to assess the environmental impacts of ammonia. Chapter 3 in: *Atmospheric Ammonia, Detecting emission changes and environmental impacts*. pp 41-48. (eds. M.A. Sutton, S. Reis and S.M.H. Baker), Springer.

Sutton M.A., van Dijk N., Levy P.E., Jones M.R., Leith I.D., Sheppard L.J., Leeson S., Tang Y.S., Stephens A., Braban C.F., Dragosits U., Howard C.M., Vieno M., Fowler D., Corbett P., Naikoo M.I., Munzi S., Ellis C.J., Chatterjee S., Steadman C.E., Möring A. and Wolseley P.A. (2020) Alkaline Air: changing perspectives on nitrogen and air pollution in an ammonia-rich world. *Phil. Trans. Roy. Soc. A* **378**: 20190315. <http://dx.doi.org/10.1098/rsta.2019.0315>

Sutton M.A., van Pul A., Sauter F., Tang Y.S and Horvath L. (2009c) Over which averaging period is the ammonia critical level most precautionary? Chapter 8 in: *Atmospheric Ammonia, Detecting emission changes and environmental impacts*. pp 93-99. (eds. M.A. Sutton, S. Reis and S.M.H. Baker), Springer.

Tang Y.S., Flechard C.R., Dämmgen U., Vidic S., Djuricic V., Mitosinkova M., Uggerud T.H., Sanz M.J., Simmons I., Dragosits U., Nemitz E., Twigg M., van Dijk N., Fauvel Y., Sanz F., Ferm M., Perrino C.,

Catrambone M., Leaver D., Braban C.F., Cape J.N., Heal M.R. and Sutton M.A. (2020) Pan-European rural atmospheric monitoring network shows dominance of NH₃ gas and NH₄NO₃ aerosol in inorganic pollution load. *Atmospheric Chemistry and Physics* **21**, 875-914. doi.org/10.5194/acp-21-875-2021

Tang, Y. S., Braban, C. F., Dragosits, U., Dore, A. J., Simmons, I., van Dijk, N., Poskitt, J., Dos Santos Pereira, G., Keenan, P. O., Conolly, C., Vincent, K., Smith, R. I., Heal, M. R., and Sutton, M. A. (2018) Drivers for spatial, temporal and long-term trends in atmospheric ammonia and ammonium in the UK, *Atmos. Chem. Phys.* **18**, 705-733. doi.org/10.5194/acp-18-705-2018.

UNECE (2007) Review of the 1999 Gothenburg Protocol. Report on the Workshop on Atmospheric Ammonia: Detecting Emission Changes and Environmental Impacts. ECE/EB.AIR/WG.5/2007/3. See: <https://digitallibrary.un.org/record/592931?ln=en> (last accessed 29 June 2022). Also available at CEH (2006).

UNECE (2017) Manual for Modelling and Mapping Critical Loads & Levels. Chapter 3: Mapping critical levels for vegetation. Coordinating Centre for Effects, Umweltbundesamt, Germany. <https://www.umweltbundesamt.de/en/manual-for-modelling-mapping-critical-loads-levels?parent=68093> (last accessed 29 June 2022).

Van Damme M., Clarisse L., Franco B., Sutton M.A., Erisman J.W., Wichink Kruit R., van Zanten M., Whitburn S., Jadj-Lazaro J., Hurtmans D., Clerbaux C. and Coheur P.-F. (2021) Global, regional and national trends of atmospheric ammonia derived from a decadal (2008-2018) satellite record. *Environmental Research Letters* **16** 055017. doi.org/10.1088/1748-9326/abd5e0

Van Damme M., Clarisse L., Whitburn S., Hadji-Lazaro J., Hurtmans D., Clerbaux C. and Coheur P.F. (2018) Industrial and agricultural ammonia point sources exposed. *Nature* **564**, 99–103.

Van Pul A., Hertel O., Geels C., Dore A.J., Vieno M., van Jaarsveld H.A., Bertström R., Schaap M. and Fagerli H. (2009) Modelling the atmospheric transport and deposition of ammonia at a national and regional scale. Chapter 19, in: *Atmospheric Ammonia: Detecting emission changes and environmental impacts* (eds. M.A. Sutton, S. Reis and S.M.H. Baker), pp 301-358, Springer.

Van Zanten M.C., R.J. Wichink Kruit, R. Hoogerbrugge, E. Van der Swaluw, W.A.J. van Pul (2017) Trends in ammonia measurements in the Netherlands over the period 1993–2014. *Atmospheric Environment*, **148**, 352-360.

Vogt E., Dragosits U., Braban C.F., Theobald M.R., Dore A.J., van Dijk N., Tang Y.S., McDonald C., Murray S., Rees R.M. and Sutton M.A. (2013) Heterogeneity of atmospheric ammonia at the landscape scale and consequences for environmental impact assessment. *Environmental Pollution* **179**, 120-131.

Wichink Kruit R.J., Aben J., de Vries W., Sauter F., van der Swaluw E., van Zanten M.C. and van Pul W.A.J. (2017) Modelling trends in ammonia in the Netherlands over the period 1990–2014. *Atmospheric Environment*, **154**, 20–30. DOI: 10.1016/j.atmosenv.2017.01.031.

Wolseley P.A., Leith I.D., Sheppard L.J., Lewis J.E.J., Crittenden P. and Sutton M.A. (2014) *Guide to using a lichen based index to nitrogen air quality*. Field Studies Council. 8 pp [ISBN 978 1 908819 10 9]. Available at: https://www.apis.ac.uk/sites/default/files/downloads/A4-Guide%20to%20the%20lichen%20based%20nitrogen%20air%20quality%20index_0.pdf (last checked 29 June 2022).

4.1.2 Literature review on the effects of ammonia

Jürgen Franzaring¹, Julia Kössler¹

¹University of Hohenheim, Institute of Landscape and Plant Ecology, Stuttgart, Germany

This presentation summarized the main findings of a background document that was prepared before the workshop and was sent to the participants. It focussed on early and more recent ammonia research and concluded that during recent years not many experiments with regard to the effect of ammonia on plants had been conducted compared to earlier years. The update of the review is presented in chapter 2 of the present document and includes several additions and modifications that were communicated to us after the workshop.

4.1.3 Modelling ammonia concentrations, trends and scenarios in Europe

Hilde Fagerli¹, Jan Eiof Jonson¹, Agnes Nyiri¹, Svetlana Tsyro¹, Wenche Aas²

¹Norwegian Meteorological Institute (EMEP/MSC-West), Oslo, Norway

²Norwegian Institute for Air Research (NILU) (EMEP-CCC), Kjeller, Norway

Whilst NH₃ emissions in Europe has been reduced only by ca 12% during the 2000-2019 period, much larger emission reductions have been reported for SO_x (ca 80%) and NO_x (ca 50%) emissions. Using the regional scale EMEP/MSC-W model, we have analyzed how these emission changes affected concentrations and wet depositions of reduced nitrogen during this period (Aas et al. 2021). Furthermore, we have studied the effects of reductions in NH₃ emissions on PM_{2.5} concentrations and depositions of nitrogen in Europe in the light of present (2017), past (2005), and future (2030) conditions (Jonson et al. 2022).

Few observational sites show statistically significant trends in wet deposition of reduced nitrogen in model results or EMEP observations for the 2000-2019 period. On the other hand, total ammonium in air (NH₃+NH₄⁺) decreased by about 25% in observations and model calculations for the same period, which is larger than the reduction in ammonia emissions. The reduction for ammonium aerosols is even larger, around 50%, both in observations and model calculations. These large differences between the changes found for the different reduced nitrogen components can be explained by the interaction of ammonia with the sulfur and oxidized nitrogen components. Furthermore, the results imply that the contribution of ammonia emissions to aerosols has been largely reduced during the 2000-2019 period, due to the impact of SO_x and NO_x emission reductions.

Assuming that the NEC Directive will be met by 2030, we calculate that reductions in PM_{2.5} levels per gram of NH₃ emissions mitigated are about a factor of 2.6 lower starting from the level of 2030 emissions compared to using 2005 emissions (due to the less efficient formation of NH₄⁺ when SO_x and NO_x emissions have been reduced much more than NH₃ emissions). Thus, mitigation of NH₃ emissions are expected to reduce PM_{2.5} less efficiently than it used to.

Following the expected reductions of NH₃ emission (19 % reductions in 2030 compared to 2005), depositions of reduced nitrogen will also decrease in Europe. However, as the reductions in NO_x emission are larger than for NH₃, the fraction of total nitrogen (reduced plus oxidized nitrogen) deposited as reduced nitrogen is increasing, and may exceed 60% in most of Europe by 2030. Thus, the potential for future reductions in the exceedances of critical loads for eutrophication in Europe will mainly rely on the ability to reduce NH₃ emissions.

A full description of these studies can be found in Aas et al. (2021) and Jonson et al. (2022).

References:

Aas, W.; Fagerli, H.; Yttri, K. E.; Tsyro, S.; Solberg, S.; Simpson, D.; Gliß, J.; Mortier, A.; Wærsted, E. G.; Brenna, H.; Hjellbrekke, A.; Griesfeller, J.; Nyíri, A.; Gauss, M.; Scheuschner, T. (2021): Trends in observations and EMEP MSC-W model calculations 2000-2019. EMEP Report 1/2021, Ch. 4

Jonson, J. E.; Fagerli, H.; Scheuschner, T. and Tsyro, S. (2022): Modelling changes in secondary inorganic aerosol formation and nitrogen deposition in Europe from 2005 to 2030. In *Atmospheric Chemistry and Physics* (22), pp. 1311–1331. DOI: 10.5194/acp-22-1311-2022

4.1.4 Measuring ammonia in the Netherlands

Susanna Rutledge-Jonker¹, Albert Bleeker¹

¹National Institute of Public Health and Environment (RIVM), Bilthoven, the Netherlands

Why do we monitor?

The National Institute of Public Health and Environment (RIVM) in the Netherlands is responsible for different environmental monitoring networks. For air quality the two relevant networks for ammonia measurements are the National Monitoring Network Air Quality (LML) and the network Monitoring Ammonia in Nature Areas (MAN).

The overall measurement objectives for the ammonia measurements are:

- Model validation and calibration
- Determining trends in time
- Performing process studies
- Reference function for other related measurements

In the next sections, a short overview is given of the different ammonia measurements in the Netherlands.

Intensive monitoring of ammonia concentrations

There are six high-accuracy ammonia measurement sites within the LML network (Figure 8). By means of miniDOAS systems (Berkhout et al., 2017), where DOAS stands for Dual Optical Absorption Spectroscopy, hourly concentrations of ammonia are collected. The miniDOAS is an open path optical system, with no tubing involved. This has the advantage that there is no delay due to NH₃ sticking to inlet lines or air filters. Furthermore, such an optical system has no interference with ammonia salts. Information on these measurements can be found at <https://www.rivm.nl/landelijk-meetnet-luchtkwaliteit> (in Dutch).

Low-cost monitoring of ammonia concentrations

The low-cost monitoring of ammonia concentrations in the Natura2000 network in the Netherlands consists of more than 500 samplers at more than 300 locations distributed over the country (Lolkema et al., 2015; Noordijk et al., 2020; Figure 8). These locations are within 86 Natura 2000 areas. The measurements are performed by means of Gradko passive samplers. Monthly concentrations are collected for about 2-9 measurement locations per Natura 2000 area, with at least one triplicate per area.

The overall advantages of the low-cost passive samplers are:

- They are small
- They are cheap
- No housing and/or electricity needed
- Monthly sampler change is easy (so samplers can be replaced by local assistants, enabling a large spatial distribution)


However, there is also an important disadvantage: the measurements are less accurate than high-accuracy instruments.

To largely overcome this accuracy issue, a hybrid design is used in the Netherlands. In this design, low-cost samplers and high-accuracy data from the LML stations are combined allowing a calibration correction on a monthly basis.

Information on the low-cost measurements in nature areas, including the monitoring results per Natura 2000 area, is available via <https://man.rivm.nl> (in Dutch).

Figure 8: Spatial distribution of intensive high-resolution miniDOAS (left) and low-cost Gradko (right) measurements in the Netherlands.

Black are current measurement locations, in red and blue the proposed locations for expansions.



Source: Wichink Kruit et al. (2021).

Dry deposition monitoring of ammonia

In the Netherlands, a small network exists for monitoring of dry deposition of ammonia. By means of a gradient technique, the dry deposition is determined by combining the concentration difference between two heights and a measure of turbulence. At four locations in the Netherlands these measurements are performed using a COTAG system (Conditional Time Averaged Gradient, e.g. Famulari et al., 2010), providing monthly depositions. Information on these measurements, including the monitoring results, can be found at www.rivm.nl/stikstof/meten/drogedepositieNH3 (in Dutch).

Research is ongoing with respect to another gradient system. Two miniDOAS systems are used in a gradient setting, providing half-hourly dry deposition values (Swart et al., in preparation).

Development in the Dutch networks

In the coming year(s) different expansions of the measurements are scheduled in the Netherlands (Wichink Kruit et al., 2021; in Dutch):

- The number of intensive miniDOAS systems will be increased from six to eight, with the two additional stations located in low concentration areas near the Dutch coast (Figure 8).

- The number of low-cost Gradko measurements will be expanded by a total of 19 locations; with ten agricultural locations and nine urban locations and at country borders (Figure 8).
- The COTAG dry deposition network will be expanded from four to ten locations, spatially distributed over the Netherlands and covering different ecosystem types.
- Furthermore, the measurements of ammonium and nitrate aerosols in the air and wet deposition of reduced and oxidized nitrogen will be expanded.
- Not only the 'traditional' monitoring activities are expanded, also the use of satellite data will be explored.

References

Berkhout, A. J. C.; Swart, D. P. J.; Volten, H.; Gast, L. F. L.; Haaima, M.; Verboom, H. et al. (2017): Replacing the AMOR with the miniDOAS in the ammonia monitoring network in the Netherlands. In *Atmospheric Measurement Techniques* 10 (11), pp. 4099–4120. DOI: 10.5194/amt-10-4099-2017.

Famulari, D.; Fowler, D.; Nemitz, E.; Hargreaves, K. J.; Storeton-West, R. L.; Rutherford, G. et al. (2010): Development of a low-cost system for measuring conditional time-averaged gradients of SO₂ and NH₃. In *Environmental monitoring and assessment* 161 (1-4), pp. 11–27. DOI: 10.1007/s10661-008-0723-6.

Lolkema, D. E.; Noordijk, H.; Stolk, A. P.; Hoogerbrugge, R.; van Zanten, M. C.; van Pul, W. A. J. (2015): The Measuring Ammonia in Nature (MAN) network in the Netherlands. In *Biogeosciences* 12 (16), pp. 5133–5142. DOI: 10.5194/bg-12-5133-2015.

Noordijk, H.; Braam, M.; Rutledge-Jonker, S.; Hoogerbrugge, R.; Stolk, A. P.; van Pul, W. A. J. (2020): Performance of the MAN ammonia monitoring network in the Netherlands. In *Atmospheric Environment* 228. DOI: 10.1016/j.atmosenv.2020.117400.

Swart, D. et al., in preparation. Measuring dry deposition of ammonia using flux-gradient and eddy covariance methods with two novel open-path instruments.

Wichink Kruit, R. et al., 2021. Op weg naar een optimale meetstrategie voor stikstof, Rijksinstituut voor Volksgezondheid en Milieu, Bilthoven.

4.1.5 Continuous monitoring of ammonia concentrations with Cavity Ring-Down Spectroscopy

Ruthger van Zwieten¹

¹Picarro B.V., Eindhoven, The Netherlands (rvzwieten@picarro.com)

High-precision trace gas measurements are essential for a range of industrial and air quality monitoring purposes. Picarro's Cavity Ring-Down analyzers (CRDS) have become the reference technology for continuous, high precision trace gas measurements, and they are used for numerous gas monitoring purposes like measuring continuously the ammonia concentration in ambient air, including *in situ* studies (Twigg et al. 2022). Key features of Picarro's CRDS analyzers are: (i) low calibration requirements, (ii) field deployable, (iii) negligible interference ('interference-free'), (iv) long term unattended operation, and (v) the possibility to measure multiple species. Here we present the performance and use cases for the Picarro G2103 NH₃ analyzer (e.g. Kamp et al. 2019) as well as the recently released Picarro G2509 multi species analyzer that allows to measure NH₃, CH₄, CO₂, N₂O and H₂O simultaneously. For NH₃, these analyzers reach a precision in the ppt range.

References

Kamp, J. N.; Chowdhury, A.; Adamsen, A. P. S.; Feilberg, A. (2019): Negligible influence of livestock contaminants and sampling system on ammonia measurements with cavity ring-down spectroscopy. In *Atmospheric Measurement Techniques* (12), pp. 2837–2850. DOI: 10.5194/amt-2018-377.

Twigg, M. M.; Berkhout, A. J. C.; Cowan, N.; Crunaire, S.; Dammers, E.; Ebert, V. et al. (2022): In-situ measurements of NH₃: instrument performance and applicability. In *Atmospheric Measurement Techniques* [preprint]. DOI: 10.5194/amt-2022-107.

4.1.6 Ammonia monitoring at long-term and temporary sites in Flanders

Jeroen Staelens¹

¹Flanders Environment Agency, Belgium

Flanders Environment Agency (VMM) has been monitoring ambient ammonia (NH₃) since 2002. The network currently consists of 17 sites located in nature areas and rural areas closer to known NH₃ sources. The air concentrations are determined using diffusive samplers and reported according to EN 17346 (2020). The main aims of the measurements are to assess annual mean NH₃ levels, to examine time trends and to calibrate and validate results of atmospheric dispersion modelling.

In 2020, the annual mean NH₃ was lower than the critical level for higher plants (3 µg/m³) at only 4 of 17 sites. The critical level for lichens and bryophytes (1 µg/m³) was not reached. Since 2008, on the average no decrease in the measured annual site means has been observed. To estimate NH₃ at the regional level, the Flemish version of the Operational Priority Substances model is applied (Sauter et al. 2018) at a grid cell resolution of 1 km². The modelled annual NH₃ is calibrated based on the measurements using a single linear regression approach.

Flanders Environment Agency also carries out temporary measurement campaigns. In 2015-2016, NH₃ was measured at 106 extra sites in nature areas. This study indicated that the permanent monitoring sites were spatially representative for the background NH₃ concentrations in Natura 2000 sites in Flanders (VMM 2017). The model slightly overestimated the measurements, but there was a good linear correspondence between the model and measurements results. In 2017, NH₃ was measured at 60 temporary sites in 6 nature areas to examine local variations. This study showed that the modelling results could be improved by combining the regional OPS model outcomes with local modelling near sources by the Immission Frequency Dispersion Model (IFDM).

Because of the important contribution of NH₃ to nitrogen deposition and secondary particulate matter, there has been increasing interest in this pollutant. In this respect, VMM investigated the possibilities of more innovative techniques such as differential optical absorption spectroscopy (DOAS) and the COnditional Time Averaged Gradient (COTAG) method. The DOAS method is an open-path technique that allows real-time measurements of NH₃ at high temporal resolution without sampling artefacts. In collaboration with RIVM we have been using two miniDOAS monitors since 2017. Recently we also set up two COTAG devices at heathland sites to determine the net dry deposition flux of NH₃, in collaboration with UKCEH.

EN 17346 (2020). Ambient air - Standard method for the determination of the concentration of ammonia using diffusive samplers.

Sauter F., van Zanten M., van der Swaluw E., Aben J., de Leeuw F., van Jaarsveld H. (2018). The OPS-model - description of OPS 4.5.2. National Institute for Public Health and the Environment (RIVM), the Netherlands.

VMM (2017). Tijdelijk meetnet ammoniak in Natura 2000-gebieden – eindrapport. Vlaamse Milieumaatschappij, Belgium. <https://www.vmm.be/publicaties/tijdelijk-meetnet-ammoniak-in-natura-2000-gebieden-eindrapport>.

4.1.7 Monitoring atmospheric ammonia on Natura2000 sites in the Republic of Ireland

David B. Kelleghan¹, Enda Hayes², Mark Everard³, Thomas P. Curran¹

¹ School of Biosystems & Food Engineering, University College Dublin, Belfield, Dublin, Ireland

² Air Quality Management Resource Centre, University of the West of England, Bristol, UK

³ International Water Security Network, University of the West of England, Bristol, UK

The Republic of Ireland has exceeded its ammonia (NH₃) emission ceiling under the National Emission Ceilings (NEC) Directive (2016/2284/EU) (European Union 2016) since 2016 (Environmental Protection Agency 2021). Additionally, Ireland’s national ambient ammonia monitoring carried out in 2013 – 2014 across 26 sites only identified three below the critical level of 1 µg NH₃ m⁻³ (Doyle et al. 2017). It was hence deemed appropriate to carry out targeted atmospheric ammonia monitoring on Natura 2000 sites protected under the Habitats Directive (92/43/EEC) (European Economic Committee 1992). From August 2017 through to July 2018, atmospheric ammonia was monitored across 12 Natura 2000 sites in the Republic of Ireland using ALPHA samplers in triplicate (Kelleghan et al. 2021a). A subset of Natura 2000 sites containing potentially sensitive qualifying interests were prioritised. Table 3 lists habitats included alongside the range of monitored atmospheric concentrations and modelled total nitrogen deposition rates for each habitat type. This work monitored airborne concentrations of 0.47 – 4.59 µg NH₃ m⁻³ across all sites. These monitored concentrations were used to calculate the dry deposition of NH₃ based on habitat specific deposition velocities (de Kluizenaar and Farrell 2000). Other forms of deposited nitrogen were extracted from EMEP modelling (The Norwegian Meteorological institute 2019) and summed to estimate total nitrogen deposited.

Table 3: Summary of atmospheric ammonia (NH₃) monitoring and modelled nitrogen (N) deposition across 12 Natura 2000 sites during 2017-2018 in the Republic of Ireland

No. of sites	Qualifying Feature	NH ₃ concentration / µg NH ₃ m ⁻³	N Deposition / Kg ha year ⁻¹
6	Active raised bogs*	1.18 – 2.92	10.1 – 17.32
1	Blanket bogs (active)	0.57	9.52
1	Transition mires and quaking bogs	1.40	11.07
2	Semi-natural dry grasslands and scrubland facies on calcareous substrates (<i>Festuco-Brometalia</i>) (* important orchid sites)	0.47 – 2.29	5.93 - 12
2	Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles	2.7 – 3.1	13.64 - 15.27
1	Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp.	4.59	17.78

Considering Active raised bogs* are a priority Annex I habitat within the Habitats Directive (92/43/EEC) and the sensitivity of *Sphagnum* moss to atmospheric ammonia, the recorded exceedance of both their critical level (1 µg NH₃ m⁻³) and empirical critical load (5 – 10 Kg ha year⁻¹) is cause for concern. Recent national modelling has also shown that modelled NH₃ concentrations across all active raised bog SACs in Ireland exceed their critical level (Bealey et al. in press). Where the Blanket bogs (active) site does not exceed its critical level for NH₃, when other forms of nitrogen are added, it exceeds the lower range of its empirical critical load (5 – 10 Kg ha year⁻¹), highlighting

the need to consider both critical levels and loads during assessment of impacts. While one Semi-natural dry grasslands and scrubland facies on calcareous substrates (*Festuco-Brometalia*) (* important orchid sites site fell below both its critical level and load, the other exceeded its critical level and its vegetation community change point (8.3 Kg ha year⁻¹) (Wilkins et al. 2016). Both relevant critical levels and loads were exceeded in all remaining SACs. Monitoring identified strong statistical relationships between monitored NH₃ and both the proportion of agricultural land within 2 km (Pearson correlation co-efficient of 0.8) and temperature (Pearson correlation co-efficient of 0.9). The relationship between temperature and ammonia concentration is potentially confounded by seasonally variable agricultural practices. This monitoring also showed a positive correlation with the MARSH (Mapping Ammonia Risk on Sensitive Habitats) model (Kelleghan et al. 2019) with a Pearson correlation co-efficient of 0.7 (Kelleghan et al. 2020). Kelleghan et al. (2020) clarified that, on at least one site (Raheenmore Bog SAC with an annual average concentration of 2.3 µg NH₃ m⁻³) ecological indicators of atmospheric NH₃ pollution were observed. Such indicators are displayed in Figure 9 including, proliferation of nitrophytic species (*Xanthoria parietina* and algae) and breakdown and decay of both *Sphagnum* sp. and *Cladonia portentosa*. This site was in receipt of NH₃ primarily from local cattle production and fertiliser/slurry application (Kelleghan et al. 2020).

Figure 9 Ecological indicators of ammonia pollution on peatland site in the Republic of Ireland



Source: Fotos from David B. Kelleghan

The National Ecosystem Monitoring Network (NEMN) is being developed in the Republic of Ireland as a response to the updated NEC Directive (Kelleghan et al. 2021b), which requires monitoring of air quality impacts across sensitive habitats. Such networks are required by the European Commission to be representative, risk-based and cost-effective. Linking with and utilising monitoring carried out by other networks is key to the success of the NEMN. Additionally, adoption of a tiered monitoring design adapted from ICP Forests monitoring minimises the cost of such a network while maximising its benefits. This design requires numerous Level 1 sites (15 minimum per habitat) where only non-invasive periodic monitoring occurs. This monitoring is dependent on other networks, such as that carried out under Article 17 of the Habitats Directive (92/43/EEC) every six years or ICP Forests crown condition surveys every four years.

The addition of parameters such as soil sampling and percent nitrogen tissue analysis (of moss species) are currently being explored, alongside inclusion of permanent quadrats (within Article 17 monitoring). Level 2 sites require some level of continuous passive atmospheric monitoring. NEMN proposes splitting these into Level 2 and Level 2 core, where Level 2 sites focus primarily on NH₃

monitoring and wet deposition where required and Level 2 core sites are intended to receive a full suite of atmospheric monitoring. This would include use of passive samplers for NH₃ and NO_x, bulk precipitation and active denuders for gases and aerosols; additional monitoring can be added as the network develops. The current NEMN design aims to have at least two Level 2 core sites per habitat, expanded by Level 2 sites to increase coverage across the risk gradient. As this is a tiered design, it is intended that each Level 2 core site will also receive Level 2 monitoring, and each Level 2 site will also function as a Level 1 site. This tiered approach ensures that relationships between atmospheric ammonia and nitrogen deposition with ecological responses can be quantified. This is an iterative network, intended to grow from year to year to monitor impacts long-term impacts. Data collected will be key to assessing and developing habitat-specific critical levels and loads.

Researchers in the University College Dublin School of Biosystems and Food Engineering have been communicating the effects of atmospheric NH₃ to the general public in Ireland through television, radio and newspaper. Their most recent endeavour is a comic suitable for children about an anthropomorphic cloud of ammonia gas. “There’s Something about Ammonia” (Wright et al. 2021) gives a basic introduction to sources, impacts and solutions to NH₃. This work has thus far been translated to 15 languages thanks to the assistance of the EPA, the COST-Action LivAge and a number of volunteers. This comic is freely available to download and use (<https://www.ucd.ie/ammonian2k/publications/comic/>).

References

Bealey, B.; Kelleghan, D. B.; Cummins, T.; Rowe, E. C.; Curran, T.; Vieno, M.; Tomlinson, S.; Carnell, E.; Sawicka, K.; Hernandez, C. M.; In press. Determination of Critical Loads for Ireland. EPA Report, Johnstown Castle, Wexford, Ireland.

Environmental Protection Agency (2021): Ireland’s Air Pollutant Emissions 1990 – 2030. Environmental Protection Agency . EPA Report, Johnstown Castle, Wexford, Ireland. https://www.epa.ie/publications/monitoring--assessment/climate-change/air-emissions/EPA-Irelands-Air-Pollutant-Emissions-report_2021Final.pdf.

de Kluizenaar, Y.; Farrell, E. P. (2000): Ammonia Monitoring In Ireland: A Full Year Of Ammonia Monitoring. EPA Research Report, Johnstown Castle, Wexford, Ireland.

Doyle, B.; Cummins, T.; Augustenborg, C.; Aherne, J. (2017): Ambient Atmospheric Ammonia in Ireland, 2013-2014. EPA Research Report, Johnstown Castle, Wexford, Ireland.

European Economic Committee (1992): Council directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. Official Journal of the European Union, 206, 22.7.1992, p. 7–50.

Kelleghan, D. B.; Hayes, E. T.; Everard, M.; Curran, T. P. (2020): Assessment of the impact of ammonia emissions from intensive agriculture installations on SACs and SPAs, EPA Research Report. Wexford, Ireland.

Kelleghan, D. B.; Hayes, E. T.; Everard, M.; Curran, T. P. (2019): Mapping ammonia risk on sensitive habitats in Ireland. In *Science of the Total Environment*, 649, 1580–1589. DOI: 10.1016/J.SCITOTENV.2018.08.424

Kelleghan, D. B.; Hayes, E. T.; Everard, M.; Keating, P.; Lesniak-Podsiadlo, A.; Curran, T.P. (2021a): Atmospheric ammonia and nitrogen deposition on Irish Natura 2000 sites: Implications for Irish agriculture. In *Atmospheric Environment*, 261, 118611. DOI: 10.1016/j.atmosenv.2021.118611

Kelleghan, D. B.; Tang, Y. S.; Rowe, E. C.; Jones, L.; Curran, T. P.; McHugh, K.; Smart, S.; Martin Hernandez, C.; Taylor, P.; Coyle, M. (2021b): National Ecosystem Monitoring Network (NEMN)-Design: monitoring air pollution impacts across sensitive ecosystems. EPA Report, Johnstown Castle, Wexford, Ireland.

The Norwegian Meteorological institute (2019): *EMEP MSC-W modelled air concentrations and depositions [WWW Document]*. Available: https://emep.int/mscw/mscw_moddata.html. Last accessed 30/03/2022.

European Union (2016): Directive (EU) 2016/2284 of the European parliament and of the council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC. Official Journal of the European Union, 344, 1–3.

Wilkins, K.; Aherne, J.; Bleasdale, A. (2016): Vegetation community change in Atlantic oak woodlands along a nitrogen deposition gradient. In *Environmental Pollution*, 216, 115–124. DOI: 10.1016/j.envpol.2016.05.024

Wright, N. T.; Kelleghan, D. B.; Curran, T.P. (2021): There's Something About Ammonia. Available: <https://www.ucd.ie/ammonian2k/publications/comic/>. Last accessed 30/03/2021.

4.1.8 National ammonia monitoring network in the UK: its current use and future potential

Khalid Aazem¹

¹ Joint Nature Conservation Committee, UK

The Department for Environment, Farming and Rural Affairs (Defra) supports work to monitor concentrations of gaseous ammonia (NH₃) and particulate ammonium (NH₄⁺) in the UK. The UK Centre for Ecology and Hydrology (UKCEH) manages the National Ammonia Monitoring Network (NAMN) for Defra.

The NAMN was established in 1996 with the objective of quantifying temporal and spatial changes in air concentrations and deposition of NH₃ and NH₄⁺ (included since 1999) on a long-term basis⁴. The monitoring provides a baseline to verify compliance with international agreements, such as the UNECE Gothenburg Protocol 1999 and the EU National Emissions Ceilings Directive NECD 2016/2284 (superseded in the UK by the National Emissions Ceilings Regulations 2018 since EU exit).

The National Ammonia Monitoring Network (NAMN) is part of the UK Air Pollution Impacts on Ecosystem Networks (APIENs). UK APIENs was formed in 2018 by integrating national air quality and ecosystem monitoring networks and surveys to meet monitoring (Article 9) and reporting (Article 10) obligations under the EU National Emissions Ceilings Directive (NECD 2016/2284). Since the UK exited the EU reporting has been transposed under Part 5 of the UK National Emissions Ceilings Regulations 2018 and is aligned to the 4 yearly European reporting cycle. APIENs is designed to monitor and report the negative impacts of air pollution (e.g., acidification, eutrophication, ozone damage and changes in biodiversity) on ecosystems that are representative of freshwater, natural and semi-natural habitats and forests in the UK⁵.

The UK ammonia monitoring network has sites located across representative areas across the UK (Figure 10). There are 52 ALPHA[®] and 27 DELTA[®] monitoring sites (Table 4) collecting spatial and temporal changes in ammonia and ammonium concentrations and deposition on a long-term basis. It is important to note that the DELTA[®] samplers also collect Nitric acid (HNO₃), Sulphur dioxide (SO₂), Hydrogen chloride (HCl), Nitrate (NO₃⁻), Sulphate (SO₄²⁻), Sodium (Na⁺), Calcium (Ca²⁺), Magnesium (Mg²⁺) as part of the UK Acid Gas and Aerosol Network (AGA-Net)⁶.

The integrated data collected will provide evidence to determine the state of the ecosystem and provide a baseline against which to measure any changes and potential recovery in ecosystem responses to emissions reductions.

⁴ UKCEH Ammonia Network <http://www.pollutantdeposition.ceh.ac.uk/content/ammonia-network> [accessed 23/05/2022]

⁵ APIENs Home page <http://www.apis.ac.uk/apiens> [accessed 12/05/2022]

⁶ Defra UK Air Information Resource: UK Acid Gas and Aerosol Network <https://uk-air.defra.gov.uk/networks/network-info?view=aganet#:~:text=The%20network%20provides%20a%20long,of%20acidity%20and%20nutrient%20nitrogen> [accessed 24/05/2022] AGA-Net is managed by the Environment Agency on behalf of Defra.

Figure 10 Distribution of UK National Ammonia Monitoring Network sites



Source: CEH Edinburgh


Table 4: The distribution of ammonia network sites by UK country (source: UKCEH)

	ALPHA (NH ₃)	27 DELTA (NH ₃ and NH ₄ ⁺)
England	37	13
Northern Ireland	1	2
Scotland	12	9
Wales	2	3
Total	52	27

Rowe et al. (2021) reported that in 2017 for 64.7% of the UK land area ammonia concentrations exceeded the 1 µg m⁻³ critical level (set for the protection of lichens and bryophytes) compared to 63.9% in 2010 (Figure 11a). This exceedance varies spatially, with Scotland having below 20% exceedance, and England and Northern Ireland having 90% or more land area exceeding this critical level. In Northern Ireland nearly 40% of the land area exceeds the 3 µg m⁻³ critical level (set for the

protection of vascular plants), rising rapidly since 2013 (Figure 11b). England and Wales have also seen increases in the proportion of their land area exceeding the $3 \mu\text{g m}^{-3}$ critical level.

Figure 11: Percentage of UK land area with ammonia concentrations exceeding critical levels: for a) lichens and bryophytes, b) vascular plants.



Source: Rowe et al. (2021)

These trends in ammonia emissions are reflected in the air pollution inventories for each of the UK countries (National Atmospheric Emissions Inventory, 2021). The normalised trends show that ammonia is the only pollutant that is significantly higher than at the 2005 reference date, especially for Northern Ireland, England, and Wales.

The UK has reported not meeting the 2020 National Emissions Ceilings Regulations ammonia target (Defra, 2022). This then triggers a review of the UK National Air Pollution Control Programme (NAPCP) to meet the 2030 target.

In recent years there has been extensive ammonia monitoring work in Northern Ireland. In addition to the three main NAMN sites, 25 ALPHA[®] and 4 DELTA[®] sites were established by DAERA in collaboration with the Agri-Food and Biosciences Institute (AFBI) and UKCEH. The objective of this project is to validate modelled NH_3 concentrations. These sites are located to achieve representative coverage geographically across locations with both high and low ammonia background concentrations. Work is underway to incorporate these data into the NAMN in the future. Some of these sites are located close to or within designated conservation sites such as Special Areas of Conservation (SAC) or Areas of Special Scientific Interest (ASSI) (AFBI, 2021).

In addition to the NI-wide network, there is an ongoing focused regime of ammonia monitoring by DAERA NIEA in collaboration with UKCEH and Ulster Wildlife (UW) on eight Special Areas of Conservation, with between 1 and 9 ALPHA[®] monitors on each site. Wet deposition bulk-rain collectors, NO_x monitoring and vegetation analysis for foliar N levels has also been initiated on selected sites (Tang et al. 2022).

In addition to monitoring, nitrogen profiles have been developed to consider the nitrogen threats across the NI designated site network and to inform conservation management. It is hoped that this suite of work can be used to improve conservation site management through a better understanding of the ammonia and nitrogen inputs at such sites.

Pending funding, Wales plans to increase ammonia monitoring with a further 24 ALPHA[®] and three DELTA[®] sites. These will be in conjunction with local monitoring campaigns and site habitat surveys. These data will be used to confirm the background concentrations of ammonia and deposition of nutrient nitrogen. It is planned to develop nitrogen and ammonia profiles and hence sensitivity for designated conservation areas to improve conservation management. There are other examples in England where Shared Nitrogen Action Plans (SNAPs) with ammonia and nitrogen data are being used to develop conservation action plans (Natural England 2018).

The existing NAMN through the establishment of APIENs is used to provide data for the UK to report under Part 5 of the NECR. With APIENs it is intended that the integrated data can be used to monitor ecosystem recovery and input into air pollution recovery strategies. Examples have described briefly how the data from the NAMN can be used to develop strategies for controlling ammonia and help conservation management.

Acknowledgments

I would like to thank the following people for providing information used in this summary:
David Vowles, Department of Environment, Farming and Rural Affairs (Defra), UK
Áine O'Reilly, Natural Environment Division, Northern Ireland Environment Agency, UK
Simon Bareham, Radioactivity and Industry Policy, Natural Resources Wales, UK
Sim Tang, UK Centre for Ecology and Hydrology, Edinburgh, UK
Ulli Dragosits, UK Centre for Ecology and Hydrology, Edinburgh, UK

References

Agri-Food and Biosciences Institute (2021). Tackling ammonia emissions in Northern Ireland: ammonia monitoring research – fourth article in a series of seven being released by AFBI to address ammonia emissions in NI

<https://www.afbini.gov.uk/news/tackling-ammonia-emissions-northern-ireland-ammonia-monitoring-research-fourth-article-series> [accessed 18/05/2022]

Defra (2022). Emissions of air pollutants in the UK – Ammonia (NH₃).

<https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-ammonia-nh3>

National Ammonia Monitoring Network <https://uk-air.defra.gov.uk/networks/network-info?view=nh3> [accessed 12/05/2022]

National Atmospheric Emissions Inventory 2021. Air pollution inventories for England, Scotland, Wales, and Northern Ireland 2005 – 2019.

https://naei.beis.gov.uk/reports/reports?report_id=1030

National Emissions Ceilings Directive NECD 2016/2284 <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv%3AOJ.L.2016.344.01.0001.01.ENG>

National Emissions Ceilings Regulations 2018

<https://www.legislation.gov.uk/ukxi/2018/129/contents/made>

Natural England (2018). Fenn's Whixall NNR Shared Nitrogen Action Plan <https://themeressandmosses.co.uk/2018/12/05/fenns-whixall-nnr-shared-nitrogen-action-plan/> [accessed 24/05/2022]

Rowe, E.C.; Sawicka, K.; Tomlinson, S.; Levy, P.; Banin, L.F.; Martin Hernandez, C.; Fitch, A. (2021): Trends Report 2021: Trends in critical load and critical level exceedances in the UK. Report to Defra under contract AQ0849, UKCEH project 07617. https://uk-air.defra.gov.uk/library/reports?report_id=1020

Tang, Y.S.; Williams, M.; Carnell, E.J.; Stephens, A.C.M.; Iwanicka, A.; Duarte, F.; van Dijk, N.; O'Reilly, A.; McCourt, A.; McHardy, F.; Fox, T.; Dragosits, U. (2022): Atmospheric ammonia assessments on six designated sites in Northern Ireland. Year 1: June 2020 – May 2021. May 2022, under review.

UNECE Gothenburg Protocol 1999. <https://unece.org/gothenburg-protocol> [accessed 23/05/2022]

4.1.9 Ammonia concentrations measurements in Germany – an overview

Alexander Moravek¹, Markus Geupel¹

¹Umweltbundesamt, Dessau, Germany

National legal and policy situation - Current status in Germany

One of the most important regulations for air quality in Germany is the EU Air Quality Directive (AAQD) and its national implementation under the 39. Federal Immission Control Act (39. BImSchV). It includes requirements for the protection of vegetation elements and crops by setting target or critical values and long-term objectives for ozone, NO_x and SO₂. Measurements have to be carried out in the rural and rural background station categories. Despite known adverse effects of ammonia (NH₃) on the vegetation and ecosystems, there is no NH₃ regulation neither in the AAQD nor in the German federal legislation, yet, to control NH₃ concentrations. In contrast, the EU Directive on the reduction of national emissions of certain atmospheric pollutants (NEC Directive) and its national implementation, the 43. Federal Immission Control Act (43. BImSchV), set nationally binding emission reduction commitments for NH₃ in order to support the reduction of ambient NH₃ to the Critical Levels of the Geneva Air Quality Convention (Recital 8 of the NEC Directive). However, although effect monitoring (of e.g. eutrophication, acidification, nutrient balances, nitrogen deposition and ozone damage) is an essential part of the Directive and its national implementation, neither the NEC Directive itself, nor the national monitoring concept contain a regulation or a guidance on the monitoring of NH₃ concentrations.

In Germany NH₃ concentration regulations are only part of the project licensing processes. Within the air quality control policies for a license of a planned project (e.g. livestock housing), the planning institution has to proof that an additional project specific NH₃ load of 2 µg m⁻³ must not be exceeded. If a Natura2000 license is required, it has to be proofed that the new project, together with the background concentration, does not lead to an exceedance of the Critical Levels.

NH₃ concentrations in Germany – Status Quo


There is evidence that the CLTRAP Critical Levels for NH₃ are exceeded over large areas in Germany. Figure 12a shows the modelled surface NH₃ concentrations for an exemplary year. It is evident that in the hotspot regions of North-West and South Germany the Critical Level of 3 µg m⁻³ for higher plants is exceeded extensively on an annual basis. For the rest of Germany, the model indicates that the 1 µg m⁻³ Critical Level for lichens and moss is exceeded as well for most regions, with only a few areas where the annual NH₃ concentration is below that level. Comparison of the modelled NH₃ distribution with satellite observations (not shown) reveal that the model identifies the NH₃ hotspot regions well, however, a systematic underprediction in some areas of Eastern Germany is apparent. In-situ measurement of NH₃ are therefore crucial to evaluated and improve atmospheric models. At the same time, more extensive *in-situ* measurements are important for local pollution assessments.

In Germany, a national NH₃ monitoring network that focuses on ecosystem protection does not exist. However, numerous NH₃ monitoring sites were established by the federal states, predominantly within the last 15 years, although no requirement to report NH₃ concentrations exists.

Measurements are performed by different institutions and are sometimes collected within other networks. To our knowledge, NH₃ measurements are currently performed by 11 out of 16 states at

more than 93 measurements sites, predominantly in rural or remote locations. In addition, the German Environment Agency measures NH₃ at seven remote background sites across Germany. At most sites, NH₃ measurements are performed using passive samplers with a biweekly or monthly sampling interval, the network of the German Environment Agency employs denuders with a weekly sampling interval. The state of Hesse operates five sites with online measurements of NH₃ using chemiluminescence analyzers at a 30 min resolution.

Figure 12: a) Map of annual mean NH₃ surface concentrations over Germany (LOTOS-EUROS simulation for 2015; source: PINETI-3 project of the German Env. Agency). b) Annual box plot statistics of NH₃ *in-situ* measurements from 2010-2019 for regional (upper panel) and remote (lower panel) locations.



Source: Own illustration. Map (a) is based on data from the PINETI-3 project of the German Environment Agency. Data for trends (b) are taken from the air quality database of the German Environment Agency (www.env-it.de/stationen/public/open.do), which includes about 2/3 of NH₃ measurement sites in Germany.

Evaluation of the *in-situ* measurements confirms the evidence from the model (Figure 12b) that the Critical Level of 1 µg m⁻³ is exceeded regularly at most sites, even at those in remote locations. The Critical Level of 3 µg m⁻³ is exceeded occasionally, strongly depending on the region. Measurements over the last decade reveal that NH₃ concentrations have been overall slightly increasing as it has been reported for other countries in Europe as well. One reason is the relative increase of NH₃ emissions compared to those of nitrogen oxides (NO_x) and sulfur dioxide (SO₂). Since the latter two act as acids in the atmosphere, their relative depletion shifts the gas to-particulate phase equilibrium to gas phase NH₃.

Relationship between Critical Loads into Critical Levels

For the monitoring of indirect effects of NH₃, such as contribution to eutrophication and related species composition change, nitrogen deposition is evaluated against ecosystem specific Critical Loads. As nitrogen deposition flux measurements (including NH₃) are challenging to make and not yet suitable for widespread monitoring, NH₃ concentration measurements could act as a proxy with a higher spatial coverage. Using a typical fraction of NH₃ dry deposition (F_{drydep_NH3-N}) to the total nitrogen deposition ($F_{totaldep_N}$) and a typical NH₃ deposition velocity (V_{dep_NH3}) – e.g. from model

evaluations – the Critical Load (CL) can be transferred to an NH₃ concentration threshold (C_{NH3_max}):

$$C_{NH3_max} = \frac{F_{drydep_NH3-N}}{F_{totaldep_N}} \cdot \frac{CL}{V_{dep_NH3}}$$

If possible, the effective deposition velocity considering NH₃ bi-directional exchange is taken. As an example, for a nitrogen sensitive ecosystem forest with $CL = 10 \text{ kg(N) ha}^{-1} \text{ a}^{-1}$, $V_{dep_NH3} = 1.5 \text{ cm s}^{-1}$ and $F_{drydep_NH3-N}/F_{totaldep_N} = 0.3$ yields $C_{NH3_max} = 0.8 \text{ } \mu\text{g m}^{-3}$. That means that at a site with a Critical Load of $10 \text{ kg N ha}^{-1} \text{ a}^{-1}$ with the given assumptions a measured concentration of $>0.8 \text{ } \mu\text{g m}^{-3}$ would indicate an exceedance of the Critical Load value.

The given example shows that critical NH₃ concentration values [$\mu\text{g m}^{-3}$] resulting from the conversion of ecosystem specific Critical Loads [$\text{kg ha}^{-1} \text{ a}^{-1}$] are in the same range as the current NH₃ Critical Levels, which also has been shown initially by Cape et al. (2009). This finding gives another argument for the scientific validity of the Critical Levels in the given range. Also, the conversion offers a potential way to derive more ecosystem specific Critical Levels in comparison to current values for the aggregated groups of higher plants und the group of lichens and bryophytes.

Table 5: Considerations for an NH₃ regulation in the EU AAQD

Motivation & advantages	Difficulties & open questions
NH ₃ affects ecosystems and PM _{2.5} formation	NH ₃ dispersion is characterised by small scale heterogeneity
A harmonization of emission reduction and air quality control could be achieved for (like for NO _x or PM) and European air quality policies could be further aligned nature protection policies	The selection of representative measurement sites is yet to be defined. Most preferably, chosen sites would represent nitrogen sensitive ecosystems. Where appropriate, in Natura2000 areas and make use of existing monitoring stations (EMEP or AAQD); affected and non-affected ecosystems
Through multiple measurements across Europe the scientific knowledge of NH ₃ distribution, ecosystem effects and PM _{2.5} -formation would systematically be improved. This knowledge would enhance the development of effective NH ₃ reduction strategies and help the setting standards and the improvements of models	It is yet to be defined, if NH ₃ would be regulated with a target value, a limit value or a critical level or if a monitoring in a first phase could be made mandatory without defining a concentration level.

Current revision of the European Ambient Air Quality Directives: Window of opportunity

With regard to the current revision process of the European Ambient Air Quality Directives currently there is a window of opportunity to include a regulation on NH₃ monitoring in the future international framework legislation. Not making use of this option would lead to the unfavorable alternative, that NH₃, one of the most concerning air pollutants of our time, would remain unmentioned in the most important European regulation for ambient air quality. Some considerations have been put together in Table 5. Against the background of those consideration the German Environment Agency strongly supports the uptake of an NH₃ regulation to the revised AAQD.

References

39. BImSchV (2010): Neununddreißigste Verordnung zur Durchführung des Bundes-Immissionsschutzgesetzes (Verordnung über Luftqualitätsstandards und Emissionshöchst-mengen). Bundesrepublik Deutschland, vertreten durch den Bundesminister der Justiz. Berlin. https://www.gesetze-im-internet.de/bimschv_39/

43. BImSchV (2018). Dreiundvierzigste Verordnung zur Durchführung des Bundes-Immissionsschutzgesetzes (Verordnung über nationale Verpflichtungen zur Reduktion der Emissionen bestimmter Luftschadstoffe). Bundesrepublik Deutschland, vertreten durch den Bundesminister der Justiz. Berlin. https://www.gesetze-im-internet.de/bimschv_43/

AAQD (2008): Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. European Parliament, Council of the European Union. <https://eur-lex.europa.eu/legal-content/DE/ALL/?uri=celex%3A32008L0050>

Cape, J.N. et al (2009). Critical Levels for Ammonia. In: Sutton, M.A., Reis, S., Baker, S.M. (eds) Atmospheric Ammonia. Springer, Dordrecht. https://doi.org/10.1007/978-1-4020-9121-6_22

NEC Directive (2016). Directive (EU) 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC. European Union. Brussels. <https://eur-lex.europa.eu/eli/dir/2016/2284/oj>

UBA German Environment Agency (2018). PINETI-3: Modellierung atmosphärischer Stoffeinträge von 2000 bis 2015 zur Bewertung der ökosystem-spezifischen Gefährdung von Biodiversität durch Luftschadstoffe in Deutschland. Martijn Schaap, Carlijn Hendriks, Richard Kranenburg, Jeroen Kuenen, Arjo Segers, Angela Schlutow, Hans-Dieter Nagel, Anja Ritter, Sabine Banzhaf. Umweltbundesamt. Dessau-Roßlau. <https://www.umweltbundesamt.de/publikationen/pinetti-3-modellierung-atmosphaerischer>

UBA German Environment Agency (2022). Considerations on the Revision of the Air Quality Directive 2008/50 EU - Air quality projections, protection of ecosystems, assessment methods, monitoring concept. Marion Wichmann-Fiebig, Marcel Langner, Ute Dauert, Simone Richter, Bryan Hellack, Markus Geupel, Alexander Moravek, Gudrun Schütze. Umweltbundesamt. Dessau-Roßlau. <https://www.umweltbundesamt.de/publikationen/considerations-on-the-revision-of-the-air-quality>

4.1.10 Metrological aspects to support environmental ammonia monitoring

Andrea Pogány¹, Nils Lüttschwager^{1,2}, Jarvis Nwaboh¹, Olav Werhahn^{1,3}, Volker Ebert¹


¹ Physikalisch Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany

² currently with: Institut für Physikalische Chemie, Georg-August-Universität Göttingen. Tammannstraße 6, 37077 Göttingen, Germany.

³ currently on leave to: Bureau International des Poids et Mesures (BIPM), Pavillon de Breteuil, 92312 Sèvres Cedex, France

Metrology is the science of measurement, ensuring comparability, reliability and accuracy of measurement results (De Bièvre 2012). Metrological traceability is a central concept in metrology, providing the link between the measurement result and the standard (i.e., a primary realization of the unit of the measurement, typically the SI system of units) through an unbroken chain of calibrations. Metrology has been serving ambient ammonia measurements through gaseous calibration standards (van der Veen 2012), and standard measurement techniques (EN 17346:2020) (European Standard 2020). The recent EMRP project “Metrology for ammonia in ambient air” (Pogány et al. 2016) aimed to strengthen the link between the gas metrology community and field measurements. The main objectives of the project included 1) improving the quality of static and dynamic reference gas standards, 2) improving the accuracy of spectroscopic measurement techniques and 3) organizing extensive inter-comparison studies. Within this project, our aim was to convert a commercial cavity ring-down spectrometer into an instrumental optical gas standard (OGS, see Figure 13), i.e., a spectrometer capable of absolute and traceable ammonia concentration measurements without the need for calibration using gaseous standards. Besides the existing reference gas standards, this approach brings further possibilities to the operation, calibration and validation of field instrumentation. An OGS can be used to calibrate or validate other instruments using gas mixtures or ambient air, and can also be used for in-situ calibration-free gas analysis. In both cases metrological traceability is provided by the traceability of the OGS, replacing the need for traceable calibration gas mixtures. The OGS concept has been realized and demonstrated for laser-based instruments detecting H₂O (Buchholz et al. 2014), CO, NO₂, as well as HCl, the latter recently validated in a direct metrological comparison (EURAMET 2021).

Figure 13: Calibration strategies using gaseous reference standards (left) vs. an instrumental optical gas standard (right).



Source: The EMRP project “Metrology for ammonia in ambient air”

In this study we used a Picarro G2103 cavity ring-down spectrometer. The spectrometer hardware was combined with a device-external, proprietary, first principles spectral data evaluation to achieve absolute concentration measurements. Our data evaluation takes into account all relevant spectral lines of NH₃, H₂O and CO₂ in the detected spectral window using a least-squares solver by the Danish Fundamental Metrology (DFM) institute (Nielsen 2001). NH₃ concentration⁷ is calculated from the integrated NH₃ line area determined from the measured absorption spectra using the Beer-Lambert law, high accuracy line intensities of the NH₃ lines (which were measured at PTB) and measured physical properties of the gas sample. By comparing to traceable reference sensors, we found that both the temperature and pressure sensors of the spectrometer are accurate within 1 %. The wavenumber axis of the measured spectra was validated by comparing the observed positions of the absorption lines to the values listed in spectroscopic databases, deviations were found to be below 1 %. We examined temperature sensitivity of the spectrometer and the gas handling system in a climate chamber. Rapid temperature changes in the range of 10 to 30 °C ambient temperature had no measurable effect on the derived NH₃ concentrations. Response time of the spectrometer was found to be in the range of minutes (10-90 % < 1.5 minutes, 1-99 % < 30 minutes in case of steps of 50, 100 and 200 ppb⁸ in both directions). Cross-sensitivity to CO₂ was found to be negligible: a step from 0 to 440 ppm CO₂ in the gas sample led to less than 0.25 % relative change in the measured NH₃ concentration. H₂O cross-sensitivity was found to be minor: a step from 0 to 10000 ppm H₂O concentration generated only 2.4 % relative change in the measured NH₃ concentration. Although these cross-sensitivities are smaller than the uncertainty of the NH₃ concentration measurements, they might be of importance in high-precision measurements or in case of extreme H₂O or CO₂ concentrations.

Table 6: Results of the field comparison of the optical gas standard (OGS) and a mobile reference gas generator (ReGaS) (Twigg et al. 2022).

Concentration provided by the ReGaS generator /ppb	Expanded uncertainty (k=2) /ppb	Concentration measured by the OGS /ppb	Expanded uncertainty (k=2) /ppb	Normalized error
0	n/a	0.14	0.62	0.23
9.93	0.18	9.72	0.66	0.31
24.51	0.43	24.09	0.92	0.41
39.95	0.69	41.09	1.30	0.77

The OGS was compared to a traceable dynamic reference gas generator, *ReGaS* based on a permeation source, which was developed by METAS, the Swiss national metrology institute (Pascale et al., 2017). We performed the comparison in the field, inside an air-conditioned mobile laboratory and found good agreement between the two standards (see Table 6, we note that more data are reported in Twigg et al. 2022), here we report only those, where a sufficient stabilization time was given). Relative expanded uncertainties of the concentrations measured by the optical gas standard were in the range of 2-6 %, comparable to that of the ReGaS (< 3 %). In Table 6, we included the so-called normalized error, being a common measure of agreement between measurement results. If the normalized error is below 1, the two results are metrologically

⁷ amount fraction in metrology

⁸ ppb and ppm: the corresponding SI units are nmol/mol and μmol/mol, respectively

compatible, i.e. the difference between them is smaller than the uncertainty of this difference. As we see in the table, the normalized error is considerably smaller than 1 indicating good agreement between the results at all measured concentrations. We note that the applied reference gas generator ReGaS is also a new development, completely independent from our OGS, thus these results promote confidence in both standards.

Finally, our OGS participated in a field measurement campaign, together with 12 other instruments including wet chemical analyzers and several spectroscopic instruments (Twigg et al. 2022). No technical issues occurred with the OGS throughout the field campaign. Comparing the OGS to the different NH₃ instruments at various ambient concentrations revealed relative deviations of up to 20 %. More details on the possible origin of these deviations including the influence of different instrument locations, sampling line designs and states, as well as different time resolutions can be found in Twigg et al. (2022).

In summary the performance of our OGS was found to be suitable for calibration and validation experiments and its applicability under field conditions has been demonstrated. Further work is planned to elaborate a detailed operation protocol including critical maintenance of the sampling line and filters to minimize sampling artifacts and to ensure reliable and comparable NH₃ measurements in ambient air.

References

Bièvre, P. de (2012): The 2012 International Vocabulary of Metrology: “VIM”. In *Accred Qual Assur* 17 (2), pp. 231–232. DOI: 10.1007/s00769-012-0885-3.

Buchholz, B.; Böse, N.; Ebert, V. (2014): Absolute validation of a diode laser hygrometer via intercomparison with the German national primary water vapor standard. In *Journal of Applied Physics*. B 116 (4), pp. 883–899. DOI: 10.1007/s00340-014-5775-4.

European Standard EN 17346:2020 EURAMET 1498 (2021): <https://www.euramet.org/technical-committees/tc-projects/details/project/comparison-of-100-umolmol-hcl-in-n2/>

Nielsen, L. (2001): Evaluation of Measurements by the Method of Least Squares. *Algorithms for Approximation IV* (170-186).

Pogány, A.; Balslev-Harder, D.; Braban, C. F.; Cassidy, N.; Ebert, V.; Ferracci, V. et al. (2016): A metrological approach to improve accuracy and reliability of ammonia measurements in ambient air. In *Measurement Science and Technology* 27 (11). DOI: 10.1088/0957-0233/27/11/115012.

Pascale, C.; Guillevic, M.; Ackermann, A.; Leuenberger, D.; Niederhauser, B. (2017): Two generators to produce SI-traceable reference gas mixtures for reactive compounds at atmospheric levels, In: *Measurement Science and Technology*. 28 (12):124002. DOI: 10.1088/1361-6501/aa870c

Twigg, M. M.; Berkhout, A. J. C.; Cowan, N.; Crunaire, S.; Dammers, E.; Ebert, V. et al. (2022): In-situ measurements of NH₃: instrument performance and applicability. In *Atmospheric Measurement Techniques* [preprint]. DOI: 10.5194/amt-2022-107.

van der Veen, A. M. H.; van Wijk, J. I. T.; Harris, K.; Goodman, C. et al. (2021): International comparison CCQM-K117 ammonia. *Metrologia* 58 (1A):08017. DOI: 10.1088/0026-1394/58/1A/08017

4.1.11 Long-term NH₃ measurements at the TROPOS research station Melpitz using a MARGA


Laurent Poulain¹, Bastian Stieger¹, Gerald Spindler¹, Markus Wallasch², Hartmut Herrmann¹

¹Leibniz Institute for Tropospheric Research (TROPOS, Germany), Leipzig, Germany

²German Federal Environment Agency (UBA), Dessau-Roßlau, Germany

Since 2010, an hourly quantification of inorganic water-soluble PM₁₀ ions and corresponding water-soluble trace gases including NH₃ is performed using a Monitor for AeRosols and Gases in ambient Air (MARGA) at the TROPOS research station Melpitz (12°56'E, 51°32'N, 86 m a.s.l.), located about 50 km to the northeast of Leipzig, Germany. The station has been in operation since 1992 to examine the impact of atmospheric long-range transport on central European background air quality (Spindler et al. 2012; Spindler et al. 2013). The site itself is situated on a meadow and is mainly surrounded by agricultural pastures and forests. Melpitz station is part of EMEP (European Monitoring and Evaluation Programme; Level 3 station; (Aas et al. 2012)), ACTRIS (Aerosols, Clouds and Trace Gases Research Infrastructure), GAW (Global Atmosphere Watch of the World Meteorological Organization), and GUAN (German Ultrafine Aerosol Network; (Birmili et al. 2009)). During several months, ammonia measurements from MARGA were compared with offline (NH₃ mini denuder, Midefix, Radiello), as well as online (PICARRO G1103) systems (Stieger et al. 2018). This unique dataset provides us the opportunity to better understand the dynamics of ammonia over years (Figure 14a), months (Figure 14b), seasons, and during the day (Figure 14c).

Figure 14: Time series of the hourly ammonia concentration measured by a MAGRA over 10 years (a), the corresponding monthly average (b), and the seasonal diurnal patterns (c)



Source: TROPOS, Leipzig.

The MARGA deployed at Melpitz was further developed and combined with an additional ion chromatography (Compact IC) to allow quantifying low molecular-weight organic acids (such as formic, acetic, propionic, butyric, pyruvic, glycolic, oxalic, malonic, succinic, malic, glutaric, and

methanesulfonic acid) in both gas and particle phases with a 2-hour time resolution (Stieger et al. 2019). Measurements were used to investigate the partitioning of the low-molecular-weight organic acids. The results show a strong deviation of the thermodynamically expected phase partitioning of the measured low-molecular-weight organic acids (Stieger et al. 2021). Recent work emphasized that the reaction between organic acids and ammonia can lead to the formation of stable organic salts (e.g. (Schlag et al. 2017). For this reason and regarding the availability of ammonia at Melpitz, ammonia might have a significant role in the observed particle-phase enrichment by inducing the formation of organic salts species. As a consequence, it appears that ammonia is not only a key species for the formation of the secondary inorganic species like ammonium sulfate and ammonium nitrate, but it could also directly impact the organic aerosol mass concentration.

References

Aas, W.; Tsyro, S.; Bieber, E.; Bergström, R.; Ceburnis, D.; Ellermann, T. et al. (2012): Lessons learnt from the first EMEP intensive measurement periods. In *Atmos. Chem. Phys.* 12 (17), pp. 8073–8094. DOI: 10.5194/acp-12-8073-2012.

Birmili, W.; Weinhold, K.; Nordmann, S.; Wiedensohler, A.; Spindler, G.; Müller, K. et al. (2009): Atmospheric aerosol measurements in the German Ultrafine Aerosol Network (GUAN). Part 1 - soot and particle number size distribution. In *Gefahrenstoffe Reinhaltung der Luft* 69 (4), pp. 137–145.

Schlag, P.; Rubach, F.; Mentel, T. F.; Reimer, D.; Canonaco, F.; Henzing, J. S. et al. (2017): Ambient and laboratory observations of organic ammonium salts in PM₁. In *Faraday discussions* 200, pp. 331–351. DOI: 10.1039/c7fd00027h.

Spindler, G.; Gnauk, T.; Grüner, A.; Iinuma, Y.; Müller, K.; Scheinhardt, S.; Herrmann, H. (2012): Size-segregated characterization of PM₁₀ at the EMEP site Melpitz (Germany) using a five-stage impactor: a six-year study. In *Journal of Atmospheric Chemistry* 69 (2), pp. 127–157. DOI: 10.1007/s10874-012-9233-6.

Spindler, G.; Grüner, A.; Müller, K.; Schlimper, S.; Herrmann, H. (2013): Long-term size-segregated particle (PM₁₀, PM_{2.5}, PM₁) characterization study at Melpitz -- influence of air mass inflow, weather conditions and season. In *Journal of Atmospheric Chemistry* 70 (2), pp. 165–195. DOI: 10.1007/s10874-013-9263-8.

Stieger, B.; Spindler, G.; Fahlbusch, B.; Müller, K.; Grüner, A.; Poulain, L. et al. (2018): Measurements of PM₁₀ ions and trace gases with the online system MARGA at the research station Melpitz in Germany – A five-year study. In *Journal of Atmospheric Chemistry* 75 (1), pp. 33–70. DOI: 10.1007/s10874-017-9361-0.

Stieger, B.; Spindler, G.; van Pinxteren, D.; Grüner, A.; Wallasch, M.; Herrmann, H. (2019): Development of an online-coupled MARGA upgrade for the 2 h interval quantification of low-molecular-weight organic acids in the gas and particle phases. In *Atmospheric Measurement Techniques* 12 (1), pp. 281–298. DOI: 10.5194/amt-12-281-2019.

Stieger, Bastian; van Pinxteren, D.; Tilgner, A.; Spindler, G.; Poulain, L.; Grüner, A. et al. (2021): Strong Deviations from Thermodynamically Expected Phase Partitioning of Low-Molecular-Weight Organic Acids during One Year of Rural Measurements. In *ACS Earth and Space Chemistry* 5 (3), pp. 500–515. DOI: 10.1021/acsearthspacechem.0c00297.

4.1.12 NADP's ammonia monitoring network (AMoN): Long-term measurements to support deposition and critical load assessments in the United States

David Gay¹, Melissa Puchalski², Emmi Felker Quinn³

¹National Atmospheric Deposition Program (NADP), University of Wisconsin-Madison, Madison, Wisconsin USA

²US EPA/Office of Air & Radiation, Washington DC, USA

³Air Resources Division, National Park Service, Denver, USA

Since 1978, the National Atmospheric Deposition Program (NADP) has tracked the status and changes in wet deposition within precipitation across the continent. Beginning in 2010, the NADP operated the Ammonia Monitoring Network (AMoN) to provide data to evaluate long-term trends in ambient ammonia concentrations and deposition. Currently, there are 116 AMoN sites across the U.S. and Canada, and over 40,000 gaseous observations of atmospheric ammonia in our publicly available database.

This talk provided details on the operation of the AMoN, including methods, available observations, analytical results along with quality assurance information, and annual temporal and spatial NH₃ concentrations over the 10-year data record. The goal was to allow for comparison to equivalent networks across Europe to improve the global monitoring and modeling assessments. Additionally, we will provide specifics from several efforts that use this ammonia data, including US E.P.A.'s Clean Air Status and Trends Network (CASTNet) and the U.S. National Park Services (NPS). EPA uses their CASTNet observations and the AMoN results to estimate dry deposition of nitrogen species through a measurement-model fusion process (CMAQ, Community Multiscale Air Quality Modeling System). The NPS uses these wet and dry deposition estimates of nitrogen, in collaboration with the NADP's Critical Loads Atmospheric Deposition (CLAD) committee, to estimate total deposition of nitrogen in Rocky Mountain National Park (Colorado), as a comparison to critical loading of N in the Park, and to help determine the sources of this excess nitrogen.

Further Reading:

EPA Measurement-Model Fusion Dry Deposition Results:

<https://nadp.slh.wisc.edu/committees/tdep/>

NADP's Total Deposition Mapping utility:

<https://clmapper.epa.gov/>

Rocky Mountain Nitrogen Initiative:

<https://cdphe.colorado.gov/public-information/planning-and-outreach/rocky-mountain-national-park-initiative>

<https://cdphe.colorado.gov/public-information/planning-and-outreach/rocky-mountain-national-park-initiative/monitoring-and>

4.2 Workshop Day 2, 29 March 2022

4.2.1 Relationships between NH₃ in urban air and the diversity of epiphytic macrolichens

Sirkku Manninen¹, K. Jääskeläinen¹, A. Stephens², A. Iwanicka², Y. S. Tang²

¹Faculty of Biological and Environmental Sciences, University of Helsinki, Finland

²UK Centre for Ecology & Hydrology, United Kingdom

Emissions of nitrogen oxides (NO_x) from traffic are implicated in changing the diversity of epiphytic lichens with a shift to nitrophytes (Davies et al. 2017). The increasing use of three-way catalysts in cars (Borsari and de Assunção 2017) and Selective Catalytic Reduction systems in diesel vehicles (Stelwagen and Ligterink 2015) reduces the emissions of NO_x but increases ammonia (NH₃) emissions (Reche et al. 2012).

NH₃ is harmful especially to acidophytic lichens (Wolseley et al. 2009). We studied the relationship between ambient NH₃ concentration and the diversity of epiphytic lichens in Helsinki (60°10'N, 24°56'E), southern Finland, to verify the results of a previous study by Manninen (2018) which suggested that NH₃ emissions from vehicles in the area may be high enough to affect lichens.

Epiphytic macrolichens were scored on *Acer platanoides*, *Ulmus glabra*, *Quercus robur*, and *Pinus sylvestris* trunks at 20 sites in Jul-Oct 2019 based on the European Standard EN 16413:2014 (Finnish Standards Association 2014). Bark samples were collected for pH measurements and lichen thalli for analysis of total N concentration. Monthly NH₃ concentrations were measured for two years (Oct 2019 - Sep 2021) with passive samplers (ALPHA[®], Tang et al. 2001). Modelled NO₂ concentrations in 2019 were taken from a concentration map provided by the [Helsinki Region Environmental Services Authority](https://www.hsy.fi/en/air-quality-and-climate/air-quality-now/annualairqualitymap/) (<https://www.hsy.fi/en/air-quality-and-climate/air-quality-now/annualairqualitymap/>). Lichens species with eutrophication scores of 1-4 were classed as acidophytes and those with 5-9 as nitrophytes (Wirth 2010) for calculating a Lichen Diversity Value separately for each tree species per site (LDV). Corresponding values for acidophytes (LDV_A) and nitrophytes (LDV_N) were also calculated (Finnish Standards Association 2014) as was the Lichen Atmospheric Nitrogen index (L_{AN}) (Wolseley et al. 2019).

The monthly NH₃ concentrations ranged from 0.00 to 3.29 µg m⁻³ and the 2-year mean concentrations from 0.15 to 1.03 µg m⁻³. The average bark pH values were: *Pinus* 3.30, *Quercus* 4.84, *Acer* 5.35, *Ulmus* 5.65. The acidophytic *Hypogymnia physodes* was the most abundant species on *Pinus*, while nitrophytic *Parmelia sulcata* (*Quercus*) and *Phycia tenella* (*Acer*, *Ulmus*) had the highest abundances on deciduous trees across the sites. The abundance of *H. physodes* and the LDV_A were negatively correlated with NH₃ concentration, NO₂ concentration, and bark pH across deciduous trees, while the L_{AN} on *Quercus* and across deciduous trees increased both with increasing NH₃ concentration and bark pH. At the tree species level, the diversity of macrolichens on *Quercus* seemed to be the best indicator of NH₃. Overall, our results showed a decrease in the abundance of acidophytes and an increase in that of nitrophytes at NH₃ concentration below the current critical level of 1 µg m⁻³ yr⁻¹ (Cape et al. 2009). The impact of traffic-derived NH₃ emissions also appeared as an increase in the total N concentration of *P. sulcata* with increasing NH₃ concentration.

References

Borsari, V.; Assunção, J. V. de (2017): Ammonia emissions from a light-duty vehicle. In *Transportation Research Part D: Transport and Environment* 51 (9), pp. 53–61. DOI: 10.1016/j.trd.2016.12.008.

Cape, J. N.; van der Eerden, L. J.; Sheppard, L. J.; Leith, I. D.; Sutton, M. A. (2009): Evidence for changing the critical level for ammonia. In *Environmental pollution (Barking, Essex : 1987)* 157 (3), pp. 1033–1037. DOI: 10.1016/j.envpol.2008.09.049.

Davies, L.; Bates, J. W.; Bell, J. N. B.; James, P. W.; Purvis, O. W. (2007): Diversity and sensitivity of epiphytes to oxides of nitrogen in London. In *Environmental pollution (Barking, Essex : 1987)* 146 (2), pp. 299–310. DOI: 10.1016/j.envpol.2006.03.023.

Finnish Standards Association (2014): Ambient air. Biomonitoring with lichens. Assessing epiphytic lichen diversity. SFS-EN 16413.

Manninen, S. (2018): Deriving nitrogen critical levels and loads based on the responses of acidophytic lichen communities on boreal urban *Pinus sylvestris* trunks. In *Science of the Total Environment* 613-614, 751-762. DOI: 10.1016/j.scitotenv.2017.09.150

Reche, C.; Viana, M.; Pandolfi, M.; Alastuey, A.; Moreno, T.; Amato, F. et al. (2012): Urban NH₃ levels and sources in a Mediterranean environment. In *Atmospheric Environment* 57 (S1), pp. 153–164. DOI: 10.1016/j.atmosenv.2012.04.021.

Stelwagen, U.; Ligterink, N. E. (2015): NH₃ emission factors for road transport. TNO report R11005, 45 pp.

Tang, Y. S.; Cape, J. N.; Sutton, M. A. (2001): Development and types of passive samplers for monitoring atmospheric NO₂ and NH₃ concentrations. In *TheScientificWorldJournal* 1, pp. 513–529. DOI: 10.1100/tsw.2001.82.

Wirth, V. (2010): Ökologische Zeigerwerte von Flechten – Erweiterte und Aktualisierte Fassung. *Herzogia* 23, 229-248.

Wolseley, P. A.; Leith, I. D.; van Dijk N.; Sutton M.A. (2009): Macrolichens on twigs and trunks as indicators of ammonia concentrations across the UK – A practical method. In: Sutton M., Reis S., Baker S.M.H. (eds), *Atmospheric Ammonia*. Springer Science + Business media B.V., pp. 101-108.

4.2.2 Results from an ammonia fumigation and a field study using endangered nitrogen-sensitive plant species

Julia Kösler¹, Jürgen Franzaring¹

¹University of Hohenheim, Institute of Landscape and Plant Ecology, Stuttgart, Germany

Ammonia (NH₃) acts as a plant nutrient and leads to eutrophication and acidification in N-sensitive ecosystems. Research on the long-term effect of low NH₃ concentrations on higher plant species is sparse. Natural gradient studies are focussing on lower plants such as lichens and an NH₃ field fumigation system has solely been used at Whim Bog, Scotland. With a greenhouse fumigation and a field gradient approach, we aimed to provide new data for the discussion of critical levels (CLE).

Greenhouse study 2020

Our first approach was a greenhouse study conducted in 2020 in Stuttgart Hohenheim (Germany). Diluted ammonia solutions were pumped into four greenhouse chambers with a peristaltic pump where one chamber acted as a control without fumigation. The solutions were allowed to drip into plastic containers from which they gently vaporized. The vaporizing gas was distributed with fans in each chamber and the air ammonia concentrations were measured weekly with Radiello® passive samplers. Additionally, temperature and humidity were determined and the four treatments were rotated between chambers in order to minimize chamber effects. Seven plant species were chosen according to a low Ellenberg N value and their threat of extinction (*Antennaria dioica*, *Arnica montana* with two different seed origins, *Carex arenaria*, *Dianthus deltoides*, *Koeleria glauca*, *Molinia caerulea* and *Pulsatilla vulgaris*). 20 plants (one per pot) were then exposed in the chambers starting on 30 April 2020.

Watering was conducted manually with rain water and half of the pots were additionally undergoing drought stress, i.e. they were receiving around one third less water. During the season, parameters like the plant length and flower number were determined until a species-individual intermediate harvest and a final harvest was conducted at the end of the season. The aboveground biomass was cut, dried and weighed for each plant. Each species was exposed for an individual duration of time, until the advent of senescence in most plants. Temperatures over the season and throughout days did not differ much between the chambers but were generally high, reaching daily maxima of over 40 °C. Measured ammonia concentrations ranged between 4 and 18 µg m⁻³ averaged over the season. The three lowest concentrations were close (4-5 µg m⁻³) to each other and the two highest concentrations were around 8 and 18 µg m⁻³. With regard to the dry treatments, all species produced less biomass in at least one harvest. Only *P. vulgaris* without intermediate harvest were not affected by the drought, corresponding to the plant's xerophytic properties.

When looking at the different ammonia concentrations, plant reactions differed from species to species. Whereas most species revealed no clear reaction pattern, the grass *M. caerulea* showed a slight trend to increase biomass production in the well-watered plants. In the dry treatments, the opposite trend could be observed. A reason for the decreased growth could be that the higher level of NH₃ could not be assimilated by the stressed plants due to more frequent stomata closure.


Summarizing the greenhouse study, we found the fumigation set-up worked well, but planned NH₃ concentrations below 10 µg m⁻³ could not be realized and did not approach zero in the control. Furthermore, high temperatures hindered plant development, i.e. flower production in many species. With regard to the dry treatment, most species reacted similarly, with a reduced biomass

production, whereas the reaction to the different ammonia treatments varied substantially not only between species but also within the same species depending on the harvests and watering (example of *M. caerulea*).

Field study 2021

The second experiment was a field gradient study conducted at a University farm Unterer Lindenhof (ULI) in the lee of livestock stables. All five stations were chosen after a NH_3 dispersal model and at all locations, NH_3 , temperature, humidity and wind parameters were measured. Furthermore, we determined the rain quantity and chemistry (ammonium and nitrate) at all stations. Plants from the greenhouse experiment as well as new species were added. These were one perennial with 20 plants per treatment, *Dianthus gratianopolitanus* and five annuals with 10 plants per treatment, *Bromus hordeaceus*, *Erodium cicutarium*, *Iberis amara*, *Legousia speculum-veneris* and *Trifolium arvense*. Plants were exposed in pots on metal tables. Due to the rather moist conditions over most of the season 2021, no drought treatment could be realized. In contrast to the previous year, no intermediate harvest was performed. For the data analyses, we still distinguished between well-watered and dry-treated plants and plants with/without intermediate harvest to investigate potential differences in growth and carry-over effects from the previous season. Measured temperatures at the five stations were rather cool throughout the season and did not differ between the stations. The weekly determined NH_3 concentrations showed a gradient between 2 and $10 \mu\text{g m}^{-3}$ averaged over the season. Station 3 revealed slightly higher concentrations than station 4, not corresponding to the underlying dispersal model.

Figure 15: Shoot dry weight (g) of *E. cicutarium*, *B. hordeaceus* and *I. amara*



Source: Own illustration, University of Hohenheim.

Plants that had already been exposed to ammonia in the previous season, were generally producing more biomass in 2021 when no intermediate harvest was performed in 2020. On the other side,

nearly no carry-over effects with regard to biomass production could be observed. With regard to the different ammonia concentrations, species varied in their reactions. Several species showed no differences between the five stations, while others produced less biomass under higher NH₃. *M. caerulea* again showed an increased growth, but this time only the last year's dry treated plants produced more biomass at higher levels of NH₃. A reason might be that plants which produced less biomass in the previous season saved more nutrients that could be used for an increased growth in the second season.

Regarding the newly added species, we found that the perennial *D. gratianopolitanus* did not react to the different ammonia concentrations, but three out of the five annuals produced more biomass. *E. cicutarium*, *B. hordeaceus* and *I. amara* responded linearly to increasing levels of NH₃ (R² of 0.94, 0.88 and 0.82, respectively, see Figure 15). The leguminous annual *T. arvense* on the contrary tended to produce less biomass with increasing NH₃ concentrations (R² 0.68) while *L. speculum-veneris* revealed no differences between the stations.

Summary

All in all, tested plant species responded differently to the different ammonia concentrations. In both seasons, some species were not affected, while others produced more or even less biomass under higher ammonia and even within one and the same species reactions could differ. We saw that NH₃ was more beneficial in annual species and that it hindered growth in other species, especially the tested legume. Therefore, we hypothesize that legumes will be more sensitive towards ammonia, a question that should be addressed in further ammonia fumigation experiments. It remains to be discussed whether only ammonia affected the plants in the field or if stations close to the pig stable might have been affected by other gaseous compounds and/or (N-containing) particles. However, the bulk deposition at the five stations showed no differences. Based on our experiments, gradient studies with pre-grown plants are preferred over greenhouse/chamber studies and we call for further ammonia studies in order to address other plant species for the refining of critical levels in the future.

4.2.3 Towards understanding shorter-term critical levels with monthly resolution measurements

Ulli Dragosits¹

¹UK Centre for Ecology & Hydrology, United Kingdom

Elevated atmospheric ammonia (NH₃) concentrations and nitrogen (N) deposition pose a substantial threat to N-sensitive vegetation in Northern Ireland (NI). Large areas of NI experience annual mean NH₃ concentrations above the critical levels, with 80% of N-sensitive habitats exposed to concentrations >1 µg NH₃ m⁻³, and 14% of N-sensitive habitats > 3 µg NH₃ m⁻³.

Ammonia monitoring is vital for verifying national modelled annual estimates of NH₃ concentrations, to quantify the level of threat to designated sites and for understanding local sources and their spatial and temporal patterns. For the study described here, local atmospheric ammonia monitoring networks were established for eight sites of European Importance (Special Areas of Conservation, SAC), combined with vegetation assessment, including vegetation N content. Atmospheric NH₃ has been measured at monthly intervals using triplicate passive ALPHA® samplers at the following designated sites in NI, with 1-9 sampling sites per designated site (all SACs are in lowland areas surrounded by agricultural activities of different levels of intensity, except for two upland sites, as specified below):

- Ballynahone Bog (since September 2014)
- Other sites (since June 2020)
 - Curran Bog
 - Garry Bog
 - Moneygal Bog
 - Peatlands Park
 - Slieve Beagh (upland site)
 - Turmennan
 - Cuilcagh Mountain (upland site)


All sampling sites except for the two upland sites, Slieve Beagh and Cuilcagh Mountain, exceed the annual 1 µg NH₃ m⁻³ critical level, and several monitoring sites exceed the 3 µg NH₃ m⁻³ critical level across the lowland sites (four examples shown in Figure 16, top). Monthly measurements show high temporal variability, both within-year and between years, the latter highlighting, for example, different wind patterns which influence the relative importance of prevailing vs. less frequent wind directions, as well as wind speed determining the build-up or dispersion of plumes.

In line with agricultural activities, much higher NH₃ concentrations are normally observed in Northern Ireland during spring in particular (with a smaller secondary peak in early autumn), when slurry and manure applications occur. Therefore, by focusing on average annual concentrations, the underlying seasonal peaks due to local activities in the vicinity of the designated sites are often not considered in modelling assessments, which do not have a sub-annual temporal resolution. Local hotspots and gradients away from sources into designated sites are also underestimated with UK national scale modelling which quantifies grid square average concentrations that smooths out the high spatio-temporal variability at the field and farm scale which is the relevant scale for assessing local impact.

Vegetation N content (Figure 16, bottom) from five sites in Northern Ireland (including two of the above sites) has been assessed using foliar ammonium (NH_4) content (and % N for a subset), from samples taken in February (across different years). In most cases, the foliar NH_4 -N concentrations correlate well with atmospheric NH_3 concentrations, at the lowland SACs embedded in agricultural land, and can be explained as due to local intensive sources within wider agricultural activities taking place. The exception is the upland site (Cuilcagh Mountain), where relatively low atmospheric NH_3 concentrations, averaging at $0.6 \mu\text{g NH}_3 \text{ m}^{-3}$ over the year ($0.1\text{-}1.2 \mu\text{g NH}_3 \text{ m}^{-3}$ monthly averages) coincide with relatively high NH_4 in the vegetation. This is expected to be due to elevated N deposition in this remote upland landscape, and a first year of wet deposition sampling is currently in progress.

Figure 16: Average annual NH_3 concentrations at four designated nature conservation sites in Northern Ireland (top) and vegetation N content (foliar ammonium) at four sites in Northern Ireland (bottom).

Moninea Bog was sampled twice, in 2007 while a poultry farm next to the bog was operational, and in 2017, seven years after the farm had ceased operation



Source: Own illustration, CEH Edinburgh.

In conclusion, it is important to understand the temporal variability of atmospheric NH_3 and the potential impact of short-term peak concentrations, in addition to the annual average values used for comparison against critical levels. It is further important to consider the large concentration gradients away from local hotspots when assessing the impact of emission sources on sensitive species, habitats and designated sites. And finally, impacts from atmospheric nitrogen on vegetation are a complex combination of concentration-based effects (through elevated NH_3) and deposition-

based impacts, in particular wet deposition at high-precipitation sites that may be relatively remote from local emission sources.

Funding

This work was mainly funded by DAERA NIEA⁹ under a project titled “Evaluating and Mitigating N Deposition Impacts on the NI Designated Site Network”.

Contributors

UKCEH Edinburgh - Sim Y. Tang, Netty van Dijk, Meg R. Williams, Ed J. Carnell, Amy C.M. Stephens, Agata Iwanicka, Fred Duarte, Isabel N. Thomas, Duncan Harvey, Mark A. Sutton

DAERA NIEA - Áine O’Reilly, Anne McCourt, Fiona McHardy

Ulster Wildlife - Trish Fox, Andy Crory, Rory Sheehan, Simon Gray, Ronald Surgenor, Roisin Grimes

⁹ DAERA - Department of Agriculture, Environment and Rural Affairs;
NIEA - Northern Ireland Environment Agency

4.2.4 Biomonitoring of ammonia with the epiphytic lichen *Hypogymnia physodes*

Karsten Mohr¹

¹Landwirtschaftskammer Niedersachsen, Oldenburg, Germany

Abstract

The usage of epiphytic lichens as bioindicator has a long tradition in biomonitoring of air pollution. The epiphytic lichen *Hypogymnia physodes*, which was very common in the past, is relatively insensitive to acidic air. This lichen species has recently been declining significantly in many rural regions of Central Europe as a result of ammonia input, decreasing precipitation acidity and climatic changes. Between 1998 and 2020, a significant decline of *Hypogymnia physodes* was observed in a Northwest German pine forest (Sandkrug). Within this period, mean ammonia concentrations varied between 2 $\mu\text{g m}^{-3}$ and 3 $\mu\text{g m}^{-3}$ and N throughfall deposition dropped from about 30 $\text{kg ha}^{-1} \text{yr}^{-1}$ to about 20 $\text{kg yr}^{-1} \text{a}^{-1}$. At the same time, the pH of pine bark increased slightly from 4.3 to 4.5. Lichen thalli exposed to different atmospheric NH_3 concentrations higher than 2.0 $\mu\text{g m}^{-3}$ in a forest clearing of Sandkrug and in the Botanic Garden of Oldenburg between 2019 and 2021 showed clear injuries already after 12 months. In the following months, these injuries intensified with significant differences. Mean NH_3 concentrations of 1.5 $\mu\text{g m}^{-3}$ led to a moderate injury of the lichen thalli.

The studies in northwestern Germany confirm the high impact of enhanced NH_3 air pollution on *Hypogymnia physodes* and other NH_3 -sensitive lichen species and provides the possibility of NH_3 -specific biomonitoring by means of mapping.

Introduction

The usage of epiphytic lichens as bioindicators has a long tradition in biomonitoring of air pollution because they react very sensitive to air pollutants. The epiphytic lichen *Hypogymnia physodes*, which was very common in the past, is relatively insensitive to acidic air pollution (Werner 1993) and has recently been declining significantly in many rural regions of Central Europe. Enhanced air pollution of ammonia is considered to be the main driver for this decrease (deBryun et al 2009, Kolk et al. 2020, Wolseley et. al 2004). In Northwest Germany, exposure experiments were carried out to get more detailed information about the sensitivity of this lichen species towards ammonia (Mohr 2021). This paper shows further results of these experiments.


Study area

The investigations in Northwestern Germany were carried out in intensively used agricultural areas and adjacent areas in the federal state of Lower Saxony where ammonia plays an important role as air pollutant. In open land, the level of NH_3 concentrations in the air is about 3 $\mu\text{g m}^{-3}$ to 6 $\mu\text{g m}^{-3}$. Higher NH_3 concentrations are measured in the vicinity of farms, which led to a die back of exposed *Hypogymnia physodes* in a short time (Mohr 2001).

For the exposure experiments with *Hypogymnia physodes* under lower air pollution conditions the pine forest “Sandkrug” as well as the Botanical Garden in the city of “Oldenburg” were chosen. In Sandkrug, ammonia air concentrations were measured in a clearing, as well as N throughfall and bark pH were measured in pine stands since 1998. Since then, the yearly mean NH_3 concentration decreased from 3 $\mu\text{g m}^{-3}$ to 2 $\mu\text{g m}^{-3}$ and the N throughfall deposition decreased from about 30 $\text{kg ha}^{-1} \text{yr}^{-1}$ to about 20 $\text{kg yr}^{-1} \text{a}^{-1}$. The pH of pine bark initially colonized with lichens increased slightly

from 4.3 to 4.5 during this period. During the exposure periods, the 4-weekly to 6-weekly NH_3 concentrations in 2 m height varied between $1.0 \mu\text{g m}^{-3}$ and $4.7 \mu\text{g m}^{-3}$ in Sandkrug and and between $1.0 \mu\text{g m}^{-3}$ and $3.8 \mu\text{g m}^{-3}$ in Oldenburg (Figure 17).

Figure 17: NH_3 concentrations (4 – 6 monthly average) in Sandkrug (3 heights: 2m, 1.5m, 0.2m) and in Oldenburg (2m)



Source: Own Illustration, Landwirtschaftskammer Niedersachsen, Oldenburg.

Method

Following VDI (2019), vital lichen thalli of *Hypogymnia physodes* from low polluted regions in Eastern Germany were fixed on a wooden board and exposed up to 2 years, until most of the lichens showed substantial injuries. In Sandkrug, several boards were exposed in different heights, which generally leads to different NH_3 air concentrations due to NH_3 absorption of the ground vegetation (Figure 18). An influence of more favorable growth conditions at lower exposure heights could not be confirmed at natural sites of *Hypogymnia physodes*. Additionally, the equally common but less NH_3 -sensitive species *Parmelia sulcata* was exposed for comparison purposes. For evaluation, the lichen thalli were photographed at one- to several-month intervals and the degree of damage was documented in 10% increments. Simultaneously, NH_3 concentrations were measured using IVL passive samplers. More detailed information about the study methods is given in Mohr (2021).

Results

Hypogymnia physodes was very common on acid barks of trees in Sandkrug and other regions in Northwest Germany until the 1990s. Subsequently, a strong decline of *Hypogymnia physodes* and other acidophytic species (e.g. *Platismatia glauca*, *Pseudevernia furfuracea*, *Tuckermanniopsis chlorophylla*, *Usnea ssp.*) was observed. Recently, only *Hypogymnia physodes* still exists sporadically on shrubs close to the ground in Sandkrug.

Thalli of *Hypogymnia physodes* sampled from remote areas were exposed on wooden panels in a clearing of the pine forest Sandkrug. After a period of 12 months, first damages were observed (5%

to 10% degree of injury, Figure 18. After a period of 2 years, the most damaged thalli were found at an exposure height of 2 m, where a mean NH_3 concentrations of $2.1 \mu\text{g m}^{-3}$ was measured in the exposure period.

Figure 18: Exposure boards with *Hypogymnia physodes* in heights between 0.2 m and 2 m in Sandkrug




Source: Own photograph, Landwirtschaftskammer Niedersachsen, Oldenburg.

At the end of the period, almost all lichen thalli in 2 m had died off. At lower heights (1.5 m, 1 m) with mean NH_3 concentrations of $1.8 \mu\text{g m}^{-3}$, the degree of injury was even on a high level between 50% and 60%. At the lowest exposure height (0.2 m) with mean NH_3 concentrations of $1.5 \mu\text{g m}^{-3}$, the injury of *H. physodes* reached 20% on average.


Figure 20 shows the progression of the degree of injury of the exposed lichens in Oldenburg, where mean NH_3 concentrations of $2.5 \mu\text{g m}^{-3}$ were determined. The first clear damage of *Hypogymnia physodes* occurred after 12 months. In the following 3 months, the lichen thalli declined successively and eventually disappeared. In contrast, the less ammonia sensitive species *Parmelia sulca* (Mohr 2001), which had been exposed on the same panel, remained largely undamaged (Figure 21).

Figure 19: Development of injury of *H. physodes* in Sandkrug within an exposure period of 2 years at different mean NH₃ concentrations (confidence interval, medium, average, min. and max. value of injury in %)



Source: Own illustration, Landwirtschaftskammer Niedersachsen, Oldenburg.

Figure 20: Development of injury (%) of *H. physodes* within 15 months in Oldenburg (confidence interval, medium, average, min. and max. value) at mean NH₃ concentrations of 2.5 µg m⁻³



Source: Own illustration, Landwirtschaftskammer Niedersachsen, Oldenburg.

Discussion

Among epiphytic lichens, the species *Hypogymnia physodes* exhibits a relatively high sensitivity to ammonia. Although other environmental conditions, such as climate warming and decreasing precipitation acidity may also play a role, its large-scale decline is essentially the result of the large-

scale increase in NH₃ pollution (Frahm 2007, Mohr 2021). The exposure experiments in Sandkrug and Oldenburg indicate that NH₃ concentrations as low as 1.5 µg m³ lead to slight damage within 2 years. Due to the high tolerance against other pollutants and environmental effects, *Hypogymnia physodes* prove to be a suitable indicator of atmospheric ammonia pollution. Thereby, mapping of *Hypogymnia physodes* and other similarly NH₃ sensitive species (e.g. *Tuckermanniopsis chlorophylla*, *Hypocenomyce scalaris*, *Platismatia glauca*, *Pseudevernia furfuracea*, *Usnea ssp.*) can provide information about the spatial distribution of atmospheric ammonia.

Figure 21: Thalli of *Hypogymnia physodes* (left) and *Parmelia sulcata* (right) before (above) and after an exposure period of 15 months in Oldenburg



Source: Own photograph, Landwirtschaftskammer Niedersachsen, Oldenburg.

Considering the unfavorable growth conditions at the exposure boards, the results of this study agree with Sutton et al. (2022), who found a decline of *Hypogymnia physodes* and other acidophytic species at NH₃ concentrations above 1.7 µg m⁻³ based on long-term observations in the UK. The protection of other rare foliose and beard species, which were often associated together with *Hypogymnia physodes* in the past, may require even lower NH₃ ambient concentrations.

References

de Bruyn, U., Linders, H.W., Mohr, K. (2009): Epiphytische Flechten im Wandel von Immissionen und Klima. Ergebnisse einer Vergleichskartierung 1989/2007 in Nordwestdeutschland. Umweltwiss Schadst Forsch, S. 63-75.

Flechtenkundliche Arbeitsgemeinschaft Nordwestdeutschland: Webseite: http://flechten-ag-nordwest.de/bot_garten_ol.php (5.3.2021)

Frahm, J.P. (2007): Die Rolle von Ammoniak bei der eutrophierenden Wirkung von Luftschadstoffen auf Flechten. KRdL im VDI-DIN (Hrsg.): KRdL-Expertenforum: Stickstoff und die Wirkungen auf die Vegetation, S. 79-86.

v.d. Kolk, H.J., Sparrius, A., Aptroot, A. (2020): Monitoring van ammoniak met korstmossen in Friesland 2019. BLWG-rapport 26, S.1-100.

Mohr, K. (2001): Stickstoffimmissionen in Nordwestdeutschland – Untersuchungen zu den ökologischen Auswirkungen auf Kiefernforsten und Möglichkeiten der Bioindikation. Dissertation Universität Oldenburg, S.1-182. <http://oops.uni-oldenburg.de/333/1/363.pdf> (3.2.2021)

Mohr K (2021): Biomonitoring von Ammoniak mit der Blasenflechte (*Hypogymnia physodes*). Gef. Reinhaltung der Luft 81/05-06. 185-191.

Sutton, M.A., van Dijk, N., Levy, P.E., Jones, M.R., Leith, I.D., Sheppard, L.J., Leeson, S., Tang, Y.S., Stephens, A., Braban, C.F., Dragosits, U., Howard, C.M., Vienne, M., Fowler, D., Corbett, P., Naikoo, M.I., Munzi, S., Ellis, C.J., Chatterjee, S., Steadman, C.E., Möring, A., Wolseley, P.A. (2020): Alkaline air: changing perspectives on nitrogen and air pollution in an ammonia-rich world. Royal Society (Hrsg.): Philosophical Transactions of the Royal Society, Section A. (2020), <https://doi.org/10.1098/rsta.2019.0315>, (18.7.2022).

VDI 2019: VDI 3957 Blatt 21: Biologische Messverfahren zur Ermittlung und Beurteilung der Wirkung von Luftverunreinigungen (Biomonitoring); Ermittlung phytotoxischer Wirkungen von Immissionen anhand der Exposition der Blattflechte *Hypogymnia physodes*. S.: 1-20.

Werner, A. (1993): Aktives Biomonitoring mit der Flechte *Hypogymnia physodes* zur Ermittlung der Luftqualität in Hannover. Bibliotheca Lichenologica 49, S. 1-113.

Wolseley P.A., James, P.W., Purvis, O.W., Leith, I.D., Sutton, M.A.: Bioindication methods for nitrogen based on community species composition. In: Sutton MA, Pitcairn CER, Withfield CP (2004): Bioindicator and biomonitoring methods for assessing the effects of atmospheric nitrogen on statutory nature conservation sites. JNCC report No. 356, S. 75-86.

4.2.5 How can we assess the impact of project contributions of ammonia on Natura2000 sites?

Rudi Uhl¹


¹FÖA Landschaftsplanung GmbH, Trier, Germany

It is very common particularly for large infrastructure projects that they lie in or in the vicinity of Natura 2000 sites. In those cases, project contributions of nitrogen meanwhile are routinely considered within appropriate assessments under Article 6.3 of the Habitats Directive (HD). Only if adverse effects of the project on the affected site can be ruled out with certainty, a project may be permitted, otherwise an exception may only be granted under strict conditions.

Among the targets protected by the HD its Annex II lists a number of mosses. Also, lichens may be characteristic of habitats listed in annex II of the HD, thus also underlying the strict protection regime. Mosses and lichens, though, may be rather prone to damage by ammonia than by deposition of reactive nitrogen compounds in general. While different evaluation schemes for nitrogen deposition within appropriate assessments have been developed and established in the EU (Hicks et al. 2011, Whitfield & McIntosh 2014), little experience is available on the assessment of impacts by ammonia.

Figure 22 shows the assessment scheme established in Germany. To assess impacts by ammonia in a similar way, (1) critical loads may be substituted by critical levels and (2) values of background deposition may be substituted by values of ammonia concentrations in the air. Modelled values of ammonia concentrations are available at least in parts of Germany, notably in Baden-Württemberg (see Pruess, this issue). Less obvious is the answer to the next question: what might be an adequate *de minimis* criterion to distinguish relevant from irrelevant project contributions?

Figure 22: Assessment scheme of nitrogen deposition impacts within German appropriate assessments under the Habitats Directive




Source: Balla et al. (2013).

While there cannot be an exhaustive answer from the scientific point of view, it would be fallacious to think that there is a way around it: every site that is not subject to an assessment within a project permit procedure must have been found to be irrelevant by some kind of consideration, so at least implicitly a “*de minimis* value” is applied to rule out effects of nitrogen compounds.

The European Court has made very clear in numerous decisions that a precautionary approach has to be taken within appropriate assessments. Any project that may do harm to the targets of the HD, even single habitats within a protected site, must be carefully assessed. For this reason, *de minimis* criteria are much lower than the ones used under the air quality legislation.

Figure 23 shows the ratio of ammonia and NO_x depositions along a road. A typical distance where – depending on average traffic and particular emission rates – a *de minimis* criterion might apply is indicated by the red line at 330 m. Rather independent of the exact location of this line, fairly exactly half of the contributions (i.e. 50 %) may be attributed to ammonia. As the *de minimis* criterion has been set in Germany originally to assess the impacts of road projects, it makes sense to determine the amount of ammonia contained within the established *de minimis* criterion .3 kg N/ha/a also as half of this value, .15 kg N/ha/a; considering the span contained within the rounded value (.25 to just less than .35) it would amount to .1749 kg N/ha/a.

Figure 23: Ratio of road traffic contributions of ammonia- (yellow) and NO_x-depositions (green) at different distances along a road (forest)



Source: Balla et al. (BMVBS) 2013, p. 91

Source: Balla et al. (2013).

In order to (re)calculate air concentrations from depositions one has to consider deposition velocity and the purely mathematical transformation of spatial units (cm * ha vs. m³) and temporal units (a to s). 1 µg/m³ NH₃ calculates thus as $v_D \cdot 2.598 \text{ kg N/ha/a}$. With the deposition velocity that is typically applied in Germany for ammonia in forests set to 2 cm/s this results in a conversion factor of 5.2. An equivalent to the *de minimis* value of .3 kg N/ha/a to be applied for the effect of road

projects would thus be $0.03 \mu\text{g}/\text{m}^3 \text{NH}_3$ (more exactly $.03366 \mu\text{g}/\text{m}^3$); values of $0.04 \mu\text{g}/\text{m}^3$ or more would not be considered irrelevant.

This approach would give a *de minimis* value which is consistent with the well-established *de minimis* criterion for traffic borne project contributions of nitrogen deposition. More conventional work will be needed then, to agree on species specific critical levels and bagatelle values of project contributions to ammonia concentrations. In the absence of field experiments or existing expert values (both of which of course would be highly appreciated) an approach would be to assess it by careful observation of the conditions giving rise to a favourable conservation status of the species under consideration.

References

Balla, S; Uhl, R.; Schlutow, A.; Lorentz, H.; Förster, M.; Becker, C.; Müller-Pfannenstiel, K.; Lüttmann, J.; Kiebel, A.; Düring, I.; Herzog, W. (2013): Untersuchung und Bewertung von straßenverkehrsbedingten Nährstoffeinträgen in empfindliche Biotope. Bericht zum FE-Vorhaben 84.0102/2009 der Bundesanstalt für Straßenwesen. Herausgegeben vom Bundesministerium für Verkehr, Bau und Stadtentwicklung, Abteilung Straßenbau, Bonn. Bearbeitende Büros: Bosch & Partner GmbH, FÖA Landschaftsplanung GmbH, Ingenieurbüro Lohmeyer GmbH & Co KG; ÖKO-DATA GmbH, in Zusammenarbeit mit Avena & BÖF - Büro für angewandte Ökologie und Forstplanung GmbH. Im Auftrag der Bundesanstalt für Straßenwesen (BAST), Bergisch Gladbach). Forschung Straßenbau und Straßenverkehrstechnik Band 1099; BMVBS Abteilung Straßenbau, Bonn; Carl Schünemann Verlag, Bremen.

Hicks, W.K., Whitfield, C.P., Bealey, W.J., Sutton, M.A. (eds.) (2011): Nitrogen Deposition and Natura 2000: Science & practice in determining environmental impacts. COST729/Nine/ESF/CCW/JNCC/SEI Workshop Proceedings, published by COST. Available at: <http://cost729.ceh.ac.uk/n2kworkshop>

Whitfield, C. & McIntosh, N. (2014): Nitrogen Deposition and the Nature Directives: Impacts and Responses: Our Shared Experiences, JNCC Report No. 521, JNCC, Peterborough, ISSN 0963-8091. Available at: <http://jncc.defra.gov.uk/page-5954>

4.2.6 Development of a trophic assessment and management system for habitat types (Directive 92/43/EEC) considering interrelations between critical levels and critical loads

Andreas Prueß¹, Siegfried Demuth², Markus Röhl³, Rainer Gryschko⁴, Heike Puhmann⁵, Markus Bernhardt-Römermann^{6,7}, Wolfgang Wilcke⁸

and other members of the StickstoffBW working groups

¹ LUBW State Institute for Environment Baden-Württemberg

² IBL Thomas Breunig Institute for Botany and Landscape Studies, Karlsruhe

³ HfWU University of Economics and Environment Nürtingen-Geislingen

⁴ regioplus engineering company, Mainz

⁵ FVA Forest Research Institute Baden-Württemberg

⁶ Friedrich Schiller University Jena, Institute of Ecology and Evolution

⁷ German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig

⁸ KIT Karlsruhe Institute of Technology, Institute of Geography and Geoecology


Introduction

With this contribution, we suggest a straightforward approach to consider atmospheric ammonia concentrations and nitrogen deposition in the implementation of the Habitats Directive. According to the Natura 2000 Habitats Directive 92/43/EEC, the German federal states must take measures to achieve or maintain the favourable conservation status of habitat types listed in Annex I. Critical levels for ammonia and critical loads for nitrogen deposition are used to assess whether there is a need for the reduction of nitrogen inputs or whether additional input is still possible. Based on the research program StickstoffBW (2019), a habitat trophic standard for critical levels and critical loads is being developed. First estimates of habitat type critical levels are published in the national habitat handbook (BfN 2021, 2022). There are three reasons to use critical ammonia levels as a leading parameter in addition to the nitrogen deposition to implement Directive 92/43/EEC: 1. Habitat trophy management (up to 10 km from agricultural sources), 2. Indicator of the nitrogen saturation of vegetation, 3. direct effect on cryptogams (lichens and mosses) and indirect effect on ferns, flowering plants, insects and mycorrhiza. In Baden-Württemberg (BW), maps of the ammonia concentration and nitrogen deposition with a resolution of 100 m x 100 m are used as the data basis to assess the current pollution (Figure 24).

We propose "trophic levels" for the application of critical levels and critical loads according the following definition. A trophic level (TR) is the long-term target level below that a favourable conservation status concerning air-borne and soil-borne nitrogen is possible. TR's are defined for all habitat types. TR1 shows a low nitrogen supply and is typically characterized by the occurrence of highly nitrogen-sensitive plant and animal species. TR3 shows a comparatively high nitrogen supply and highly nitrogen-sensitive species are absent, but the general species composition still corresponds to a defined habitat type according to Directive 92/43/EEC. TR2 shows an intermediate nitrogen supply. The cause for high nitrogen supply can be natural (e.g. inherently high soil fertility, strong biological nitrogen fixation) or anthropogenic (e.g. former fertilization, high deposition or lack of biomass export). When setting the trophic level parameters, we assume that growth will not be limited by other factors. Growth conditions and hence biomass production typically increase from TR1 to TR3.

There is a strong interrelation between critical levels and critical loads: 32% - 49% of the nitrogen deposition in BW is caused by ammonia dry deposition (10th - 90th percentile, data from 44 BW districts; StickstoffBW 2022).

Figure 24: Ammonia concentration and nitrogen deposition in Baden-Württemberg, average for the years 2012-2016



Source: LUBW (2020).

This implies that a critical load of 20 kg N ha yr⁻¹ corresponds to a NH₃ critical level of 2.0 – 3.1 µg m⁻³ (10th - 90th percentile), when a deposition velocity v_d of 1.2 cm s⁻¹ (typical for lowland hay meadows in BW, habitat 6510) is assumed.

We will derive the habitat-specific critical level of NH₃ from the habitat-specific critical loads for nitrogen considering the trophic level with the help of the following criteria: 1.0 - 2.0 µg m⁻³ if a high proportion of very sensitive cryptogams, insects or mycorrhiza belong to the characteristic habitat species and 2.0 - 3.0 µg m⁻³ if less sensitive cryptogams, insects or mycorrhiza belong to the characteristic habitat species. 1 µg m⁻³ is taken as the field detection limit (minimum) and 3 µg m⁻³ is taken as the upper limit to protect vegetation as a nitrogen sink (maximum), also taking into account a maximum critical load of 20 kg ha yr⁻¹ and a v_d of 1.2 cm s⁻¹.

Outlook and conclusions

The future assumptions, formulas and rules for the habitat-specific critical levels and critical loads are currently being compiled in a technical guidance document (update of StickstoffBW 2019). In future, specific attention should be paid to the atmospheric ammonia concentration as an easily measured key parameter for habitat trophy management (considering critical loads). Biomass thresholds for the different trophic levels and removal of N by biomass need to be assessed more accurately. Biomass production is the most important trophic indicator. Long-term integrated field monitoring of sensitive species including cryptogams, mycorrhizae and insects (not just ferns and flowering plants) with respect to trophic levels is needed for all sensitive habitat types. Establishing a general habitat trophic standard based on critical levels and critical loads is required for the implementation of directive 92/43/EEC. Critical surplus values for agricultural land should follow soon. We consider our suggested approach as a basis for further discussion among and within research institutions, consultant offices and federal and state administrations.

References

BfN Bundesamt für Naturschutz [Ed.] (2021, 2022): Das europäische Schutzgebietssystem Natura 2000 - Naturschutz und Biologische Vielfalt NaBiV Heft 172, <https://bf.n.buchweltshop.de>, Bonn.

LUBW Landesanstalt für Umwelt Baden-Württemberg (2020): Daten- und Kartendienst der LUBW, <https://udo.lubw.baden-wuerttemberg.de/public/>

StickstoffBW (2019): Ermittlung der Critical Levels und Critical Loads für Stickstoff - Methodik für die Neufassung der Belastungsgrenzen für in Deutschland vorkommende Vegetationseinheiten (CL Bericht 2019). - LUBW Publikationsdienst ID 69710, 62 Seiten, Download <https://pd.lubw.de/69710>, Karlsruhe.

StickstoffBW (2022): Reaktiver Stickstoff in der Atmosphäre von Baden-Württemberg - Ammoniakkonzentration und Stickstoffdeposition 2012 bis 2016 (Depositionsbericht 2022).- LUBW Publikationsdienst ID 10362, 175 Seiten, Karlsruhe [in preparation].

4.2.7 Ammonia concentrations in Switzerland compared to critical levels and loads

Reto Meier¹

¹Swiss Federal Office for the Environment, Air Pollution Control and Chemicals Division, Switzerland

Measurement network

In 2020, ammonia passive sampler measurements in Switzerland and the Principality of Liechtenstein included a total of 89 sites. 13 sites have been operational since 2000, eight were added in 2004, 11 in 2008, 31 in 2013 and the remaining afterwards (Seitler 2021a; Seitler & Meier 2022). While many sites are situated in rural areas characterized by intensive livestock and crop farming, the measurement network also covers extensive alpine pastures, bogs and fens as well as traffic and urban sites. Additional measurements of ammonia and other nitrogen compounds are part of the National Air Pollution Monitoring Network (NABEL 2020). They include continuous NH₃ measurements at three EMEP sites (CRDS), daily concentration of the sum of reduced (NH₃ & NH₄⁺) as well as the sum of oxidized (HNO₃ NO₃⁻) nitrogen compounds at two EMEP sites (impregnated filters), 2-weekly concentrations of NH₃, NH₄⁺, HNO₃, NO₃⁻ at four sites (three EMEP sites, minidenuders) and wet deposited NH₄⁺ and NO₃⁻ at three sites (two EMEP sites). Further measurements are carried out by cantons (bulk deposition (Seitler 2021b), minidenuders).

Ammonia concentrations

The highest annual ammonia concentrations are found in areas with intensive animal production (6–10 µg/m³) followed by areas with predominant crop farming (2–5 µg/m³). Peak concentrations occur during periods of slurry application. Measurement sites within a few kilometres may show different concentration values, but nevertheless show a parallel pattern over time. Besides a relatively large year-to-year variability, influenced by meteorological conditions, measurements show no change in average ammonia concentrations in the last 20 years. Preliminary analysis including data on NH₄⁺ indicates, however, a slight decrease of total reduced nitrogen compounds.

Critical Load and Critical Level exceedances

On average in Switzerland, about half of the NH₃ concentration at a sensitive ecosystem is caused by sources within a radius of 4 kilometres; the other half is due to regional background (EKL 2014). In rural areas with intensive agriculture, ammonia dry deposition accounts for the major part of total nitrogen deposition. In other regions, total nitrogen deposition is largely influenced by wet deposition of NH₄⁺ and NO₃⁻. As a consequence, Critical Loads for nitrogen are more often exceeded than Critical Levels for ammonia (Rihm and Künzle 2019, BAFU 2020). The Federal Commission for Air Hygiene therefore concluded that the need for action to protect sensitive ecosystems becomes much more apparent in an assessment based on total nitrogen deposition than in an assessment based on gaseous ammonia concentrations only.

Ammonia concentration and lichens

A Swiss study (Urech et al. 2015) analysing the relationship between lichen vegetation and nitrogen pollution found that occurrence of nitrogen sensitive species steeply decreases above ammonia concentrations of 1 µg/m³. Based on lichen data over a period from 2001 to 2010 from 1815 trees in 10 different regions covering an ammonia gradient from 1 to 9 µg/m³ the authors concluded that the Critical Level for lichens needs to be substantially lower than 3 µg/m³.

References

BAFU (2020): Vollzugshilfe Übermässigkeit von Stickstoff-Einträgen und Ammoniak-Immissionen (available in German/French/Italian),

<https://www.bafu.admin.ch/bafu/de/home/themen/luft/publikationen-studien/publikationen/uebermaessigkeit-von-stickstoff-eintraegen-und-ammoniak-immissionen.html>

Federal Commission for Air Hygiene (EKL) (2014): Ammoniak-Immissionen und Stickstoffeinträge (available in German and French), https://www.ekl.admin.ch/inhalte/ekldataien/themen/Ammoniak-Immissionen_und_Stickstoffeintraege.pdf

NABEL (2020): Technischer Bericht zum Nationalen Beobachtungsnetz für Luftfremdstoffe (German), https://www.bafu.admin.ch/dam/bafu/de/dokumente/luft/fachinfo-daten/technischer_berichtzumnationalenbeobachtungsnetz fuerluftfremdsto.pdf.download.pdf/technischer_berichtzumnationalenbeobachtungsnetz fuerluftfremdsto.pdf

Rihm, B.; Künzle, T. (2019): Mapping Nitrogen Deposition 2015 for Switzerland, <https://www.bafu.admin.ch/dam/bafu/en/dokumente/luft/externe-studien-berichte/mapping-nitrogen-deposition-2015-for-switzerland.pdf.download.pdf/Mapping%20Nitrogen%20Deposition%202015%20for%20Switzerland.pdf>

Seitler, E.; Meier, M. (2021a): Ammoniakimmissionsmessungen in der Schweiz 2000 bis 2020, Messbericht, Forschungsstelle für Umweltbeobachtung (German), https://www.bafu.admin.ch/dam/bafu/de/dokumente/luft/externe-studien-berichte/NH3_Bericht_2000-2020.pdf.download.pdf/NH3_Bericht_2000-2020.pdf

Seitler, E.; Meier, M.; Ehrenmann, Z. (2021b): Atmosphärische Stickstoff-Deposition in der Schweiz 2000 bis 2019, Forschungsstelle für Umweltbeobachtung <https://www.bafu.admin.ch/dam/bafu/de/dokumente/luft/externe-studien-berichte/atmosphaerische-stickstoff-deposition-in-der-schweiz-2000-2019.pdf.download.pdf/Stickstoffdeposition2019.pdf>

Seitler, E.; Meier, M. (2022): Ammoniakimmissionsmessungen in der Schweiz 2000-2021, Messbericht, Forschungsstelle für Umweltbeobachtung FUB. Available online at https://www.bafu.admin.ch/dam/bafu/de/dokumente/luft/externe-studien-berichte/ammoniak-immissionsmessungen-in-der-schweiz-2000-2021.pdf.download.pdf/Ammoniak-Immissionsmessungen_CH_2000-2021.pdf

Urech, M.; et al. (2015): Flechtenvegetation und Stickstoffbelastung: Zeitlich und räumlich differenzierte Datenanalyse anhand epidemiologischer Methoden (German), <https://www.bafu.admin.ch/dam/bafu/de/dokumente/luft/externe-studien-berichte/flechtenvegetationundstickstoffbelastung.pdf.download.pdf/flechtenvegetationundstickstoffbelastung.pdf>

4.3 Workshop Discussion

4.3.1 Discussion points

During the workshop and the final discussion, it became clear that the discussing workshop participants see ammonia critical levels as a very important measure for ammonia mapping and ecosystem protection. It was worked out that critical levels play an equally important role as critical loads and that both approaches are instruments that need to be differentiated and looked at. Besides the critical levels for ammonia levels for NO_x might also need revision.

The workshop participants agreed that in many countries there is a lack with regard to ammonia monitoring and urge that ammonia is considered in the future review and revision of the EU Air Quality Directive. Furthermore, a Europe-wide standardization of ammonia measurements would be a useful addition to verify the effectiveness of mitigation measures needed to meet the reduction commitments of the NEC Directive.

With regard to the questions raised prior to the ammonia workshop, some answers could be collected:

- **Is the use of passive samplers suitable? Are there devices to continuously monitor NH_3 in the low range?**

The workshop participants agreed that the easy and cheap use of passive samplers is a well-established method for NH_3 measuring. It was stated that the measuring height plays an important role and that measuring should therefore always take place at a defined and standardized height (compare with European standard). Measuring ammonia in the low concentration range is generally seen as difficult, in part due to the stickiness of ammonia to instrument surfaces of real time monitors. It is recommended to monitor ammonia additionally with the denuder method or continuous online analysers in order to calibrate the passive sampler employed in the field.

It is worth noting that as a consequence of the changing partitioning between gaseous NH_3 and aerosol NH_4^+ the gaseous NH_3 trends do not necessarily reflect the changes in ammonia emissions. Additional monitoring of NH_4^+ trends (aerosol and precipitation) is important for making such a comparison (cf. Key Finding 5, chapter 4.3.2).

- **Do you perform or are you aware of research on ammonia effects using controlled fumigation?**
- **Do you perform ammonia fumigation studies in the field?**

New research using controlled fumigation is planned in Sri Lanka by UK CEH. The study site for the set-up of a fumigation system is a woodland and the impact on the site-specific vegetation shall be investigated. Furthermore, in 2021, CEH has established a new field ammonia fumigation system at Glencorse Field Facility (Southwest of Edinburgh). This site releases ammonia at a similar level to Whim, but assesses the impact of ammonia within a woodland environment. The concentration range goes up to $40 \mu\text{g m}^{-3}$.

- **Do you perform or are you aware of gradient studies using active or passive biomonitoring?**

Gradient studies mainly using passive biomonitoring with lichens are planned in Finland (S. Manninen) and in the UK (M. Jones) using e.g. Alpha samplers or CRDS analysers. In this context, the question was raised how to differentiate between oxidised/reduced N compounds adding to N deposition, i.e. how to evaluate the sole effect of ammonia on vegetation and species composition in field studies.

- **Would you support annual, monthly and/or short-term CLe?**

It was suggested that the provisional monthly CLe of 23 was too high, but new findings that support or contradict the value are missing. Generally, CLe for annual ammonia are supported due to missing sufficient new evidence to change them.

- **Would you support to include knowledge on CLo into scientific justification of CLe?**

It was recognized that there is a relationship between critical loads and levels that can be used in the assessment of NH₃ effects on vegetation and species composition change. For the conversion, assumptions on the prevailing NH₃ deposition velocity and on the ratio of NH₃ deposition to other non-NH₃ N deposition fluxes at a location need to be made (e.g. taken from model results). It was shown that critical levels from critical loads are in the same range as existing CLe between 1-3 µg and therefore give further support to the robustness of the existing CLe. All participants found it important to keep both approaches, but there need to be further discussions on whether or not to include information on the relationship between the two metrics into the Mapping Manual.

- **Is there enough evidence for changing/keeping the CLe?**

As only few new studies address the effects of ammonia concentrations on plants, all participants agreed that more studies, including higher and lower species are needed. Field fumigation studies should cover different climatic conditions and ecosystem types and the established vegetation as well as biomonitoring plants should be involved. Particular attention should be paid to important habitat-specific species.

However, based on the few new findings, the annual critical levels of 1 µg m⁻³ for lichens and bryophytes and of 3 µg m⁻³ for higher plants can be confirmed, i.e. there is no evidence to change these values. In this context, it was also stated that the long-term ammonia CLe are of higher relevance than the monthly CLe that were referred to as provisional values in the 2006 Mapping Manual. Regarding the monthly CLe of 23 µg m⁻³ there are no new findings that support or contradict these values.

- **Which further steps after the Workshop should be taken in the review of ammonia CLe?**

Workshop recommendations and conclusions, in particular for the Air Quality Directive (see key findings, chapter 4.3.2)

Further information/reading:

- EMEP data to be included (consider data exclusion rules such as data has to be gathered for at least 75% of the year, at least one month per season) is very welcome. Please contact Wenche Aas or Kjetil Tørseth at CCC/NILU for more details
- ICP Vegetation review on NO₂ Critical Levels (first online meeting 24th May 2022) → contact Mike Perring if interested
- Standard method for the determination of the concentration of ammonia using diffusive samplers: <https://www.en-standard.eu/csn-en-17346-ambient-air-standard-method-for-the-determination-of-the-concentration-of-ammonia-using-diffusive-samplers/>
- Ammonia comic in 16 languages: <https://www.ucd.ie/ammonian2k/publications/comic/>
- Papers on Gradko measurements:
<https://www.sciencedirect.com/science/article/pii/S1352231020301394?via%3Dihub>
<https://www.sciencedirect.com/science/article/pii/S1352231020302624>
- NI-wide network data will be published in <https://eidc.ac.uk/>
This is the Environmental Information Data Centre (EIDC) is part of the Natural Environment Research Council's (NERC) Environmental Data Service and is hosted by the UK Centre for Ecology & Hydrology (UKCEH). EIDC manages nationally-important datasets concerned with the terrestrial and freshwater sciences
- More information on Ireland's NEC network design:
<https://researchrepository.ucd.ie/bitstream/10197/12330/3/National%20Ecosystem%20Monitoring%20Network%20%28NEMN%29-Design.pdf>
- Paper on different passive samplers:
<https://www.sciencedirect.com/science/article/abs/pii/S1352231018308185?via%3Dihub#fig4>
- Link to report on decision making thresholds:
<https://hub.jncc.gov.uk/assets/6cce4f2e-e481-4ec2-b369-2b4026c88447>
- Research in ecotrons in V-Connecterra (University of Belgium) in Natura2000 areas:
<https://www.terhills-nationaalparkhogekepen.be/over-connecterra/veldstudiecentrum/608>

4.3.2 Key findings

The following **key messages** and **recommendations** emerged from the Dessau Ammonia Workshop:

1. The workshop participants agreed that ammonia should be better respected in the future environmental legislation. At present, there are no ammonia air quality limit values set in the European Air Quality or Habitat Directive. Given the upcoming review of the AAQD, and recognizing the inclusion of impacts on ecosystems as an essential part of both directives, it is recommended that future revisions of the AAQD directives should contain a Europe-wide standardization of ammonia measurements, to verify the effectiveness of mitigation measures needed to meet the reduction commitments. A consistent monitoring of ammonia with harmonization of data gathering and processing strategies across Europe will help to improve EU policies by laying the basis for further understanding and assessment of the effects on ecosystems, on plants and plant communities.

2. The monitoring of air pollution impacts under Article 9 (and Annex V) of the National Emissions reductions commitments Directive (NECD) offers the possibility to include NH₃ monitoring data. In contrast to several soil and vegetation parameters ammonia critical levels or air concentration measurements are not explicitly mentioned. The workshop participants concluded that the reporting of ammonia concentrations in relation to the required vegetation and soil data could be a good complement with regard to the observation of the effectiveness of the directive and that a standardized protocol should be developed in the near future.

3. Air quality limit values for human health include total PM_{2.5} and there are substantial nitrogen-related contributions to PM_{2.5} through a) reaction of NO_x emissions, b) reaction of NH₃ emissions and c) ammonia-stimulated oxidation of SO₂ emissions to SO₄²⁻ levels. The successful reductions of SO₂ and NO_x emissions slow down the formation rate of secondary inorganic PM, which is reflected in substantially reducing trends of ammonium (NH₄⁺) in PM_{2.5} in Europe and North America over the last 20 years. At the same time, ammonia residence times are increased, which is reflected in the slight upward trend in levels seen on both sides of the Atlantic.

4. European standards for passive monitoring of ammonia have been published by the Comité Européen de Normalisation (EN 17346 (2020)), and metrological comparisons of widely used passive ammonia samplers have been performed within the European MetNH₃ project. The workshop participants agree that a wide range of passive sampler implementations may achieve cost-effective monitoring of ammonia with typically monthly and bi-weekly time-integrated sampling, for the purpose of assessing long-term trends and spatial patterns. However, the performance of a method depends on specific laboratory implementation. For the setup of a monitoring network, therefore it was recommended to include parallel sampling with a recognized monitoring device on a continuous basis across the range of concentrations at selected sites, to allow ongoing calibration of passive samplers against a reference method. Currently, various real-time monitors for low concentrations of ammonia are being introduced to the market, which can be used for the calibration of passive samplers.

5. The area-wide exceeding of the compensation point due to high fertilization of croplands and grasslands and substantial emission-reductions of SO₂ and NO_x have altered the partitioning between gaseous NH₃ and aerosol NH₄⁺. For this reason, the assessment of trends in ammonia emissions and concentrations requires simultaneous measurement of the nitrogen surplus in

agriculture, gaseous NH_3 , particulate matter NH_4^+ and of NH_4^+ in rain. Several datasets show gaseous NH_3 increasing while $\text{PM}_{2.5}$ NH_4^+ is simultaneously decreasing (s. above). Higher levels of ammonia may also be due to stronger volatilization caused by rising temperatures. The workshop participants therefore strongly encourage to continue long-term studies on changes in the chemical and physical climate and their relationships to the biosphere. Future research should also make use of remote sensing technologies. The initiative of the European Space Agency to develop the NitroSat candidate mission for 2030 is greatly to be welcomed, and if finally selected, would provide unprecedented improvement in understanding the fine scale distribution and trends of ammonia and other nitrogen gases globally.

6. There have only been few long-term studies on the effects of elevated NH_3 concentrations on vegetation in the last 15 years. Existing studies covered in an extensive review (see chapter 2 for documentation) broadly support the ammonia critical levels established by the 2006 UNECE Ammonia Workshop. However, it is noted that further long-term experimental studies at a field scale for different habitat types and climates must be a priority. Under a changing pollution climate (with less SO_2 and a more alkaline atmosphere and climatic change), there is a risk that some of the most sensitive species groups (e.g. higher species, certain lichens, mycorrhiza or insects) may become more sensitive to high NH_3 concentrations. The workshop participants therefore advocate for efforts to include ammonia and nitrogen deposition in existing ecological research networks, e.g. the LTER sites. Furthermore, more basic research is needed e.g. in the NH_3 driven distortion of algae and fungi in lichens and the metabolism of ammonia/ammonium in higher plants.

7. Reduced and oxidized nitrogen forms and their adverse ecological and environmental effects cannot be disentangled from each other. While the last revision of ammonia took place in 2006, the UNECE Critical Level for NO_2 has not been reviewed for 25 years and it is unlikely to be sufficiently precautionary. Since it was originally set for “average” plant species and crops, it may not be suited to protect the most sensitive species, e.g. lichens and oligotrophic higher species. A process to review the UNECE NO_2 critical level has recently been started and will yield important information for the reduced nitrogen species.

8. The workshop confirmed the validity of annual UNECE Critical Levels (CLe) of $1 \mu\text{g m}^{-3}$ for lichens and bryophytes and of $3 \mu\text{g m}^{-3}$ for higher plants. CLe account for both direct (e.g. toxic) and indirect effects (e.g. species composition). Although species composition change is more associated with N-deposition and eutrophication, the mathematical conversion of critical nitrogen deposition values to NH_3 concentration with standardized meteorological assumptions showed a good agreement with the UNECE Critical Levels, giving further evidence for their scientific robustness.

9. Specific points should be clarified with respect to the UNECE ammonia critical levels agreed in Edinburgh in 2006, which focus on long term protection of ecosystems and sensitive organisms:

a) The 'long-term' critical levels, expressed as annual mean NH_3 concentrations of 1 and 3 [2-4] $\mu\text{g m}^{-3}$, cannot be assumed to give protection for long periods of more than 20 to 30 years.

b) The monthly critical level of $23 \mu\text{g m}^{-3}$ had been derived for higher plants, and does not apply to lichens, bryophytes and ecosystems where these are important to ecosystem integrity, e.g. peatlands. It can easily be seen that one month of $23 \mu\text{g m}^{-3}$, would give a minimum annual average of $1.9 \mu\text{g m}^{-3}$, which already exceeds the long-term critical level. To be mathematically consistent with the long-term critical level, a maximum monthly value of $12 \mu\text{g m}^{-3}$ would apply,

though further evidence would be needed to assess whether this value is sufficiently precautionary for sensitive lower and higher plant species.

10. To deal with the above-mentioned uncertainties, the workshop participants outlined the need for further basic and field-based research to refine critical levels. While ammonia field fumigation systems have been established in peatlands and forests in the UK, research infrastructures are not available in other countries and in other nitrogen-limited ecosystems, e.g. highly protected acidic and calcareous grasslands. Besides the set-up and technical improvement of ammonia fumigation systems in established vegetation, studies in the lee of livestock facilities using plants (on-site and exposed) along the ammonia gradient will create new dose-response relationships that may be used to update critical levels in the future. Studies should focus on single species as well as on species interactions, which will in the long-term lead to the disappearance of the sensitive components at the expense of the nitrogen loving species.

Meanwhile, on 4th of April 2022, these workshop recommendations have been presented by UBA at the EU Stakeholder consultations workshop on the revision of the AAQD (Policy area 3: Strengthening air quality monitoring, modelling and plans).

The note to the EU Stakeholder Consultations was as follows.

The Ammonia Expert Workshop 2022, held in the framework of the Convention on Long-Range Transboundary Air Pollution on 28 and 29 March 2022 (Dessau/online) draws the following conclusion and recommendations in relation to the revision of the EU Ambient Air Quality Directives:

Ammonia impacts ecosystems and contributes to the loss of biodiversity. In addition, it is a key precursor in the formation of PM_{2.5}. It is likely that critical levels of ammonia in the ground-level air layer, as set by Convention on Long-Range Transboundary Air Pollution (CLTRAP) and as confirmed by the review of recent data within the above mentioned 2022-workshop, are exceeded for a substantial number of ecosystems across Europe.

The workshop points out the deficiency that at present there is no obligation on ammonia monitoring and procedure for assessment set in the Ambient Air Quality Directives. Given the ongoing process to revise the AAQDs, and recognizing that impacts on ecosystems and vegetation are already now an essential part of the directive (as already reflected in ozone, NO_x, and SO₂ assessments), the workshop members urge that ammonia is considered in the future review and revision of the AAQD.

Addressing ammonia in the AAQDs would harmonize EU legislation with the NEC Directive and is essential for effective implementation of the Habitats Directive. A Europe-wide standardization of ammonia measurements, would be a useful addition to verify the effectiveness of mitigation measures needed to meet the reduction commitments of the NEC Directive. Furthermore, a consistent monitoring of ammonia with harmonization of data gathering and processing strategies across Europe will help to improve EU policies in three ways:

(1) It will help to lay the basis for further understanding and assessment of the effects on ecosystems, on plants and plant communities.

(2) Available data will help to improve chemical-transport models through evaluation and validation.

(3) and lastly the combination of both (1) and (2) will lay the basis for effective reduction strategies.

A low-cost method for the determination of the concentration of ammonia using diffusive samplers is given by the European Standard method EN 17346.

5 Acknowledgements

We are grateful for funding from the German Environment Agency (UBA). The present review and a fumigation/gradient study were part of the project: “Review of internationally proposed critical levels for ammonia” (ReFoPlan-project FKZ 3718 63 2010). The background document was sent to registered workshop participants before the meeting and an updated version is included in chapter 2 of the proceedings. We are grateful to various workshop participants who commented on the paper and made useful suggestions and additions to the document.

6 List of registered participants

Name	Institution	Country
Aazem, Khalid	Joint Nature Conservation Committee	UK
Aherne, Julian	Trent University	Canada
Alice, JAMES	ICP Modelling & Mapping / INERIS	France
Alonso, Rocío	Ecotoxicology of Air Pollution, CIEMAT	Spain
Banzhaf, Sabine	Freie Universität Berlin	Germany
Bareham, Simon	Natural Resources Wales	UK
Bastian, Susanne	Sächsisches Landesamt für Umwelt, Landwirtschaft und Geologie	Germany
Bittman, Shabtai	AAFC	Canada
Bleeker, Albert	RIVM	The Netherlands
Buckley, Máire	Environmental Protection Agency	Ireland
Buxbaum, Iris	Umweltbundesamt GmbH	Austria
Campbell, Claire	Scottish Environment Protection Agency	UK
Carrasco Molina, Tania	Centro de Investigaciones Medioambientales, energéticas y tecnológicas	Spain
Chelin, Lisa	Université Paris Est Créteil and CNRS	France
Czorny, David	LANUV NRW	Germany
Dammers, Enrico	TNO	The Netherlands
De Marco, Alessandra	ENEA	Italy
Demuth, Siegfried	Institut für Botanik und Landschaftskunde	Germany
Dennier, Silja	Drägerwerk AG Lübeck	Germany
Deventer, Julian	ANECO Institut für Umweltschutz GmbH & Co.	Germany
Diana, Pitar	National Institute for Research and Development in Forestry	Romania
Diemer, Jürgen	Bayerisches Landesamt für Umwelt	Germany
Dragosits, Ulli	UK Centre for Ecology * Hydrology	UK
Ellermann, Thomas	Aarhus University, Department for Environmental Science	Denmark
English, Yvonne	EPA, Ireland	Ireland
Fagerli, Hilde	EMEP MSC-West / Norwegian Meteorological institute	Norway
Fauvel, Yannick	INRAE	France
Ferenczi, Zita	Hungarian Meteorological Service	Hungary
Flechard, Chris	INRAE	France
Fornasier, Maria	ISPRA	Italy
Forster, Manuela	Bayer. Landesanstalt für Wald und Forstwirtschaft	Germany
Franzaring, Jürgen	University of Hohenheim	Germany
Frohn, Lise	Aarhus University	Denmark
Frößl, Jonas	University of Hohenheim	Germany
Gay, David	University of Wisconsin Madison	USA
Geels, Camilla	Aarhus University	Denmark
Gergócs, Veronika	Centre for Agricultural Research	Hungary
Geupel, Markus	German Environment Agency	Germany
Gillespie, Colin	Scottish Environment Protection Agency	UK
Götze, Hannah	Thünen Institut für Agrarklimaschutz	Germany
Grima, Alistair	Environment and Resources Authority	Malta
Gryschko, Rainer	regioplus Ingenieurgesellschaft	Germany
Hainsch, Andreas	Staatliches Gewerbeaufsichtsamt Hildesheim	Germany
Häni, Christoph	Berner Fachhochschule	Switzerland
Hayes, Felicity	ICP Vegetation	UK
Hellack, Bryan	German Environment Agency	Germany

Hellsten, Sofie	IVL Swedish Environmental Research Institute	Sweden
Hoffmann, Saskia	Niedersächsisches Minist. für Umwelt, Energie, Bauen und Klimaschutz	Germany
Hofmann, Magdalena	Picarro	The Netherlands
Höhle, Juliane	Sächsisches Landesamt für Umwelt, Landwirtschaft und Geologie	Germany
Horlacher, Dieter	KTBL	Germany
Horváth, László	Department of Optics and Quantum Electronics, University of Szeged	Hungary
Hünová, Iva	Czech hydrometeorological Institute	Czech Republic
Jazvo, Kristina	Croatian Meteorological and Hydrological Service	Croatia
Jefimow, Maciej	Institute of Environmental Protection – National Research Institute	Poland
Jenkinson, Holly	Natural England	UK
Jones, Matthew	UKCEH	UK
Junk, Jürgen	Luxembourg Institute of Science and Technology (LIST)	Luxembourg
Jutterström, Sara	CDM - IVL Swedish Environmental Research Institute	Sweden
Kaltenegger, Katrin	CIAM / IIASA	Austria
Karwisch, Stefan	HORIBA EUROPE GmbH	Germany
Kelleghan, David	University College Dublin	Ireland
Kiebel, Achim	FÖA Landschaftsplanung GmbH	Germany
Knight, Lydia	Natural England	UK
Köhler, Jutta	Bayerisches Landesamt für Umwelt	Germany
Kösler, Julia	University of Hohenheim	Germany
Kowalska, Anna	Forest Research Institute	Poland
Kozlova, Natalia	IEEP – branch of FSAC VIM	Russian Federation
Krein, Andreas	Administration de l'environnement Luxembourg	Luxembourg
Krischke, Fabian	Bayerisches Landesamt für Umwelt	Germany
Kujundzic, Olivera	Ministry of Environment	Montenegro
Kukowski, Sina	Thünen-Institut für Agrarklimaschutz	Germany
Kupper Thomas	Berner Fachhochschule Hochschule für Agrarwissenschaften HAFL	Switzerland
Langner, Marcel	German Environment Agency	Germany
Laurent, Poulain	Leibniz Institute for Tropospheric Research (TROPOS)	Germany
Lett, Céline	Luxembourg Institute of Science and Technology (LIST)	Luxembourg
Lohberger, Michael	Sächsisches Landesamt für Umwelt, Landwirtschaft und Geologie (LfULG)	Germany
Maasikmets, Marek	Estonian Environmental Research Centre	Estonia
Machon, Attila	Hungarian Meteorological Service, Air Quality Reference Centre	Hungary
Makkonen, Ulla	Finnish Meteorological Institute	Finland
Malchykhina, Hanna	Institute for Nature Management, National Academy of Science	Belarus
Manninen, Sirkku	University of Helsinki	Finland
Martin, Nick	National Physical Laboratory	UK
Mathieu-Üffing, Barbara	Lufthygienische Überwachung Schleswig-Holstein, LLUR	Germany
Meier, Mario	FUB - Forschungsstelle für Umweltbeobachtung AG	Switzerland
Meier, Reto	Federal Office for the Environment	Switzerland
Meremans, Dominique	Flemish environmental agency	Belgium
Mohr, Karsten	Chamber of Agriculture in Lower Saxony	Germany
Moravek, Alexander	German Environment Agency	Germany
Nagl, Christian	Umweltbundesamt	Austria
Nauber, Andreas	Drägerwerk AG Lübeck	Germany
Niebuhr, Jana	German Environment Agency	Germany
O'Reilly, Aine	Northern Ireland Environment Agency	UK
Pacholski, Andreas	Thünen Institut für Agrarklimaschutz	Germany
Perring, Mike	UKCEH	UK
Philippe, Maetz	Belgian Interregional Environment Agency	Belgium
Pogány, Andrea	PTB (Physikalisch-Technische Bundesanstalt)	Germany

Pompe, Frank	UMW Umweltmonitoring	Germany
Prescher, Anne	ICP Forests	Germany
Pröll, Gisela	Environment Agency Austria	Austria
Prüß, Andreas	LUBW Baden-Württemberg	Germany
Puchalski, Melissa	EPA Office of Air & Radiation, CASTNET Program Manager	USA
Pulis, Kristian	Environment & Resources Authority	Malta
Rábago, Isaura	WGE -CIEMAT	Spain
Richter, Simone	German Environment Agency	Germany
Rüdiger, Julian	UBA	Germany
Rutledge-Jonker, Susanna	RIVM	The Netherlands
Sanz Noriega, Carolin	UNECE	UNECE
Saul, Melanie	Thünen Institut für Agrarklimaschutz	Germany
Schaap, Martijn	TNO	The Netherlands
Schmidt, Ulrike	Staatsbetrieb Sachsenforst	Germany
Schmitz, Andreas	LANUV	Germany
Schmoeckel, Gerhard	Bayerisches Landesamt für Umwelt	Germany
Schrader, Frederik	Thünen Institute of Climate-Smart Agriculture	Germany
Seitler, Eva	FUB - Forschungsstelle für Umweltbeobachtung AG	Switzerland
Sharps, Katrina	UK Centre for Ecology & Hydrology	UK
Spangl, Wolfgang	Umweltbundesamt	Austria
Stack, Philip	Environmental Protection Agency	Ireland
Staelens, Jeroen	Flanders Environment Agency	Belgium
Steiner, Anne	German Environment Agency	Germany
Stevens, Carly	Lancaster University	UK
Sutton, Mark	TFRN / UK Centre for Ecology & Hydrology (UKCEH)	UK
Tang, Sim	UKCEH	UK
Tatzber, Michael	Federal Research and Training Centre for Forests (BFW)	Austria
Tiggemann, Luzia	FÖA Landschaftsplanung GmbH	Germany
Trebs, Ivonne	Luxembourg Institute of Science and Technology (LIST)	Luxembourg
Twigg, Marsailidh	UK Centre for Ecology & Hydrology	UK
Uhl, Rudolf	FÖA Landschaftsplanung GmbH	Germany
Valach, Alex	Applied University of Bern	Switzerland
van Dijk, Netty	UKCEH Edinburgh	UK
van Zwieten, Ruthger	Ruthger van Zwieten	Switzerland
Vermeulen, Simon	Walloon Agency for Air and Climate	Belgium
Verstraeten, Arne	Research Institute for Nature and Forest (INBO)	Belgium
Wäber, Monica	UMW Umweltmonitoring	Germany
Wall, Brendan	Environmental Protection Agency	Ireland
Watne, Ågot	IVL Swedish Environmental Research Institute	Sweden
Werhahn, Olav	PTB	Germany
Windisch, Ute	Technische Hochschule Mittelhessen	Germany
Yakimova, Elena	Ministry of Environment and Water	Bulgaria
Zhang, Jun	EMSA, TNO	The Netherlands